

PEEL HERE

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.
Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility.
If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Harmful if inhaled. May irritate the nose, throat and skin. May irritate the eyes. Avoid inhaling spray mist. Avoid contact with eyes and skin.
When opening the container and preparing spray, wear elbow-length chemical resistant gloves. After each days use, wash gloves and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126).

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any state and Federal Legislation or law of a Territory. Not for repacking or reformulations. No license under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 80777/101615

Batch No:

Date of Manufacture:

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE
---	--



AXICHEM PTY LTD
A.B.N 35 131 628 594
9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

**DIRECTIONS FOR USE
FOR ALL STATES**

A. APPLIED AS AN IN-FURROW TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400mL	4 weeks (grazing)
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	200 or 400mL	
	Septoria tritici blotch (<i>S. tritici</i>) Take-All (<i>Geaumannomyces graminis</i>)	400mL	
Barley	Powdery Mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	200 or 400mL	

CRITICAL COMMENTS

Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.

Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide.

Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)
60	667*
70	571
80	500
100	400
120	333

For the control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.

Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide.

Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	
	200mL rate	400mL rate
50	400	800*
60	333	667*
70	286	571
80	250 ⁺	500
100	200 ⁺	400
120	167 ⁺	333

*Application volumes over 600mL per 100kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.

⁺ Where application volume is low, product may be diluted with water to improve coverage.

B. APPLIED AS FOLIAR SPRAY

Wheat	Stripe rust (<i>Puccinia striiformis</i>)	250-500mL Ground application: Apply in 100-150L water plus 200mL of a 1000g/L non-ionic surfactant per 100L of spray solution. Aerial application Apply in a minimum of 15L water plus 200mL a 1000g/L non-ionic surfactant per 100L of spray solution.	7 weeks (Harvest or grazing)
	Leaf rust (<i>P. recondita</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>), Septoria tritici blotch (<i>S. tritici</i>)		
Barley	Powdery mildew (<i>Erysiphe graminis</i>)		10 weeks (harvest or grazing)

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

A. AC FLOUT 250 SC APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. AC FLOUT 250 SC APPLIED AS A FOLIAR SPRAY

BARLEY - DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION

- DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 10 WEEKS OF TREATMENT.

WHEAT - DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION

- DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 WEEKS OF TREATMENT.

GENERAL INSTRUCTIONS

Fertiliser application

Ensure the treated fertiliser is dry before sowing the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

Foliar Applications

Check the crop carefully and regularly. Spray only if there is sufficient time for the rust to develop to a level on the variety in question when it will cause significant yield loss.

GROUP 3 FUNGICIDE

Fungicide Resistance Warning

AC Flout 250 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to AC Flout 250 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by AC Flout 250 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of AC Flout 250 SC Fungicide to control resistant fungi.

Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

Apply after the flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

Protection of Wildlife, Fish, Crustaceans and Environment

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

Storage and Disposal

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Safety Directions

Harmful if swallowed. Harmful if inhaled. May irritate the nose, throat and skin. May irritate the eyes. Avoid inhaling spray mist. Avoid contact with eyes and skin.

When opening the container and preparing spray, wear elbow-length chemical resistant gloves. After each days use, wash gloves and contaminated clothing.

Wash hands after use.

First Aid

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126).

Safety Data Sheet

For further information refer to the Safety Data Sheet which is available from the supplier.

LABEL ELEMENTS

GHS Label Elements	
SIGNAL WORD	WARNING
Hazard Statement(s)	
H302	Harmful if swallowed.
H312	Harmful in contact with skin.
H319	Causes serious eye irritation.
H332	Harmful if inhaled.
H412	Harmful to aquatic life with long lasting effects.
Precautionary statement(s) Prevention	
P271	Use only outdoor or in a well-ventilated area.
P261	Avoid breathing mist/vapours/spray.
P264	Wash all exposed external body areas thoroughly after handling.
P270	Do not eat, drink or smoke when using this product.
P273	Avoid release to the environment.
P280	Wear protective gloves, protective clothing, eye protection and face protection.

Precautionary statement(s) Response

P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+P313	If eye irritation persists: Get medical advice/attention.
P301+P312	IF SWALLOWED: Call a POISON CENTRE/doctor/physician/first aider if you feel unwell.
P302+P352	IF ON SKIN: Wash with plenty of water.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P330	Rinse mouth.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

Not Applicable.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 80777/101615

AxiChem