


PULL HERE

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**ACTIVE CONSTITUENT: 250 g/L AMITROLE
220g/L AMMONIUM THIOCYANATE**

GROUP Q HERBICIDE

For the control of weeds in orchards, vineyards, irrigation ditches and drains, roadsides, wheat and barley, and for general industrial situations.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

STORAGE AND DISPOSAL

Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. **DO NOT** burn empty containers or product.

Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 80927/101955

Batch No:

Date of Manufacture:

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE
---	--

AxiChem
Pty Ltd

AXICHEM PTY LTD
A.B.N 35 131 628 594
9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

DIRECTIONS FOR USE

SITUATION AND CROP	WEEDS CONTROLLED	STATE	Rate/Ha
Wheat and Barley	Rye Grass, Wild Oats	All States	2.8L 5.6L
Vineyards and Orchards	Couch Grass		1.1L/100L water
	Broadleaf Weeds and Grasses		570mL/100L water
Potatoes Pre-Harvest Preparation	Rye Grass, Winter Grass, Capeweed, Paterson's Curse	Vic, Tas only	5.5-11L
<i>Pinus radiata</i> Plantations	Rye Grass, Wild Oats, Yorkshire Fog Grass, Silver Grass, Sand Brome, Capeweed, Clovers, Mouse-ear Chickweed, Sow Thistle and Spear Thistle seedlings	NSW, SA, WA, Tas, Vic only	5.6-8L/ha plus 2.8-4L/ha flowable Atrazine 500g/L
Non-crop areas around buildings, Commercial and Industrial areas, Domestic and Public Service areas, Right-Of-Ways	Blackberry	Vic, Tas, SA only	2L/100L water
	Furze or Gorse, Ox-eye Daisy		1.1L/100L
	Fennel, Hoary Cress, African Daisy, Apple of Sodom, Illyrian Thistle		
Aquatic Areas, Drains and Channels, Margins of Streams, Lakes and Dams	Russian (creeping) Knapweed, Watsonia, Artichoke Thistle, Prairie Ground Cherry, Onion Weed, Khaki Weed, Chilean Cestrum (Green Cestrum), Oxalis (Soursob), Couch, Kikuyu, Fog Grass, Sorrel	All States	
	Cumbungi, Phragmites and Nutgrass		2.3L/100L water
	Water Couch		4.5L/100L water 2.2L/100L water
	Water Hyacinth		280mL/100L water

NOT TO BE USES FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

ORCHARDS AND VINES: DO NOT APPLY LATER THAN 56 DAYS BEFORE HARVEST

OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED

CRITICAL COMMENTS
Pre-planting. Sow the crop at a depth of 5cm not less than 5 days after spraying. A mechanical cultivation 4 to 5 days after spraying will assist results.
Apply when grasses are in the 2-3 leaf stage and 5cm high.
Apply when grasses are in the 6 leaf stage and 10cm high.
DO NOT use on sandy soils in cropping situations.
Apply when weeds are small and actively growing. Repeat application in 6-8 weeks. Apply as a directed spray to weeds only.
Apply approximately 4-5 weeks before harvest after the potato haulms have dried off. Use the higher rates when rye grass is in clumps or other weeds more than 30cm high are present. For boom spraying apply at least 100litres of spray mixture per hectare. If other broadleaf weeds such as Clovers, Variegated Thistle and Cruciferous weeds are also present add a Low Volatile Ester 400 Herbicide. Warning: if used on sandy soils do not plant follow-up crops within 3 months.
Use highest rates where grasses are a problem. Spray evenly. Pre-planting: Graze area heavily and apply the tank mixture not more than 2 weeks before planting.
Post-planting: Apply tank mixture immediately after or within one month of planting. Do not spray pine seedlings using directed spray.
Apply twice. First treatment in December or January and regrowth sprayed in autumn. Respraying in the following year is essential. DO NOT apply to blackberry bushes carrying mature or developing fruit.
Apply during summer or before full flowering. Respraying will be necessary.
Apply when weeds are actively growing, immediately prior to flowering.
Apply when weeds are actively growing, immediately prior to flowering. Respraying will be necessary to destroy regrowth and seedlings.
Apply during flowering period between January and May. Re-treatment of Nutgrass will be necessary.
Apply as a single treatment between February and May.
Apply in November to December with a further application at the same rate 6-8 weeks later.
Apply immediately prior to flowering.

GENERAL INSTRUCTIONS

Before opening carefully read Safety Directions, First Aid Instructions, Directions for Use, Storage and Disposal and Protection Statements.

RESISTANT WEEDS WARNING

GROUP Q HERBICIDE

AC Amon Herbicide is a member of the triazole group of herbicides. AC Amon Herbicide has the inhibitors of carotenoid biosynthesis mode of action. For weed resistance management AC Amon Herbicide is a Group Q herbicide. Some naturally occurring weed biotypes resistant to AC Amon Herbicide and other Group Q herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by AC Amon Herbicide or other Group Q herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Amon Herbicide to control resistant weeds. AC Amon Herbicide may be subject to specific resistance strategies. For further information contact your farm chemical supplier, consultant, local Department of Agriculture or local Axichem Pty Ltd representative.

APPLICATION INFORMATION

Thoroughly wet all surfaces by aiming to apply 1700-2200 Litres per hectare with power spraying equipment, 900-1100 Litres per hectare (1 Litre per 10m²) using a knapsack and 170 Litres per hectare when using a misting machine.

Note: Misting machines are not recommended for use in Tasmania.

MIXING INSTRUCTIONS

To improve the efficiency of this product it is recommended that a 700g/L active surfactant be added to the spray mixture at the following rates:

Power Spraying – 250mL/100L spray mixture

Knapsack Spraying – 55mL/15L knapsack

Misting Machine – 60mL/15L spray mixture (Not recommended in Tasmania)

Note: The rates indicated in the Directions for Use table are per 100L of water for power spraying. Refer to the chart below for the required amounts to suit other equipment.

CONVERSION CHART FOR RECOMMENDATIONS TO SUIT VARIOUS EQUIPMENT

Recommended Rate per 100L of water	Per 15L of knapsack	Per 5L Misting
4.5L	1.1L	1.3L
2.2L	570mL	630mL
2.0L	510mL	560mL
1.7L	430mL	480mL
1.1L	280mL	310mL
570mL	110mL	120mL
280mL	55mL	60mL

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: This product is a non-selective herbicide and may damage crops, pastures and some ornamentals. However it is safer to use near plants susceptible to phenoxy type herbicides. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Equipment Maintenance and Usage:

Do not use the spray machine for any other purpose unless thoroughly cleaned by several rinsings of all parts with clean water.

STORAGE AND DISPOSAL

Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Safety Directions

DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

First Aid

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Safety Data Sheet

For further information refer to the Safety Data Sheet which is available from the supplier.

LABEL ELEMENTS

GHS Label Elements	
---------------------------	--

SIGNAL WORD	DANGER
--------------------	---------------

Hazard Statement(s)

H312	Harmful in contact with skin.
H332	Harmful if inhaled.
H319	Causes serious eye irritation.
H350	May cause cancer.
H361d	Suspected of damaging the unborn child.
H373	May cause damage to organs through prolonged or repeated exposure.
H411	Toxic to aquatic life with long lasting effects.
AUH032	Contact with acid liberates very toxic gas.
H302	Harmful if swallowed.

Precautionary statement(s) Prevention

P201	Obtain special instructions before use.
P260	Do not breathe mist/vapours/spray.
P271	Use only outdoors or in a well-ventilated area.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P264	Wash all exposed external body areas thoroughly after handling.
P270	Do not eat, drink or smoke when using this product.
P273	Avoid release to the environment.

Precautionary statement(s) Response

P308+P313	If exposed or concerned: Get medical advice/attention.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+P313	If eye irritation persists: Get medical advice/attention.
P391	Collect spillage.

P301+P312	IF SWALLOWED: Call a POISON CENTRE/doctor/physician/first aider if you feel unwell.
P302+P352	IF ON SKIN: Wash with plenty of water.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P330	Rinse mouth.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

P405	Store locked up.
-------------	------------------

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

AxiChem