

PEEL  
HERE

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENTS: 150 g/L TRIADIMENOL  
4 g/L CYPERMETHRIN

**GROUP 3 FUNGICIDE**      **GROUP 3A INSECTICIDE**

Wheat: For control of bunt, flag smut and loose smut and suppression of seedling stripe rust and speckled leaf blotch.  
Barley: For control of covered smut, loose smut and powdery mildew and suppression of leaf scald.  
Oats: For control of loose and covered smut.  
For protection against insect pests of stored grain.

## IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

### STORAGE AND DISPOSAL

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product. DO NOT allow seed treated with this product to contaminate grain intended for human consumption. DO NOT use seed treated with this product for human consumption. Bags which have held treated seed are not to be used for any other purpose.

**For Non-Refillable Containers:** Store product in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse empty containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For Refillable Containers:** Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs AxiChem Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

May irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

### LIMIT OF WARRANTY AND LIABILITY

AxiChem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of AxiChem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 81969/104511

Batch Number:

Date of Manufacture:

<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	<b>FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE</b>
---	--

**AxiChem**  
Pty Ltd

**AXICHEM PTY LTD**  
A.B.N 35 131 628 594  
9 PALINGS COURT, NERANG QLD 4211  
PHONE: 07 5596 1736 www.axichem.com.au

**AxiChem**

**DIRECTIONS FOR USE:****RESTRAINTS:** DO NOT treat physically damaged seed.

CROP	DISEASE/PEST	RATE	CRITICAL COMMENTS
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut. Suppression of seedling infections of Stripe Rust.	100 mL/ 100 kg seed	AC Haven Flowable Seed Dressing should be mixed with water to give even coverage of seed. The quantity of water used for mixing will vary depending on type of equipment and quality of seed. Use a minimum of 400 mL of mixture (i.e. AC Haven Flowable Seed Dressing plus water) with each 100 kg of seed. DO NOT use more than 600 mL of mixture (product + water) with each 100 kg of seed. Whatever dilution is used it is essential that 100 mL (or 150 mL) of AC Haven Flowable Seed Dressing is applied/100 kg seed. The mixture should be gently stirred regularly. If nasal irritation occurs wear a facemask while treating seed. Use the 150 mL/100 kg seed rate where foliar diseases frequently occur and where any of the following are required: - longer suppression of Stripe Rust on seedlings - control of seedling infections of Powdery Mildew - suppression of Speckled Leaf Blotch in wheat - extended suppression of Leaf Scald in barley: and use higher rate on more susceptible varieties or where disease pressure is heavy.
Oats	Covered and Loose Smut.		
Barley	Covered and Loose Smut. Suppression of seedling infections of Powdery Mildew.  Suppression of seedling infections of Leaf Scald (not Queensland).		
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut. Suppression of seedling infections of Stripe Rust (not Queensland).  Suppression of seedling infections of Speckled Leaf Blotch (not Qld and WA).	150 mL/ 100 kg seed	Both rates give protection against insect pests. Use rate appropriate to disease control required.
Barley	Covered and Loose Smut. Control of seedling infections of Powdery Mildew. Suppression of seedling infections of Leaf Scald (not Queensland).		
Wheat Barley Oats	Protection against insect pests of stored grains: Granary Weevil, Indian Meal Moth, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle, Tropical Warehouse Moth.	100 or 150 mL/ 100 kg seed	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:****DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 5 WEEKS OF SOWING.****GENERAL INSTRUCTIONS****In Wheat**

AC Haven Flowable Seed Dressing, especially at the higher rate of 150 mL/100 kg seed, may delay emergence. Crops will recover rapidly in most soil types. However, seed treated with the 150 mL/100 kg seed rate should not be sown in soils that are likely to crust or clod at the surface after sowing. Sowing AC Haven Flowable Seed

Dressing treated wheat deeper than 50 mm may slightly reduce percentage plant establishment in heavier soils and sowing rates may need to be adjusted accordingly. The possible delay in emergence of wheat treated with the 150 mL/100 kg seed rate may counteract the advantage of suppression of Speckled Leaf Blotch.

### **Storage of treated seed**

Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment may be necessary. If the seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Although AC Haven Flowable Seed Dressing has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for the performance of stored treated seed.

### **FUNGICIDE RESISTANCE WARNING**

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

AC Haven Flowable Seed Dressing is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

### **INSECTICIDE RESISTANCE WARNING**

<b>GROUP</b>	<b>3A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management AC Haven Flowable Seed Dressing is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to this product and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if this product or other Group 3A insecticides are used repeatedly. The effectiveness of this product on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects. AC Haven Flowable Seed Dressing may be subject to specific resistance management strategies. For further information contact your local supplier, Axichem representative or local agricultural department agronomist.

### **SEED QUALITY**

AC Haven Flowable Seed Dressing seed treatment should not be used on seed with more than 12% moisture content, or on sprung, sprouted, damaged or severely pinched seed or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard. Treating with AC Haven Flowable Seed Dressing as directed will raise the moisture level by up to 0.6%, depending on weather conditions at treatment. The use of AC Haven Flowable Seed Dressing mixed with water at recommended rates will have no effect on the storage life of treated sound seed.

### **CLEANING UP**

Equipment should be thoroughly cleaned with water. In the event of changing from this water based flowable formulation to a solvent-based liquid formulation, it is essential to rinse equipment with methylated spirits

after thoroughly cleaning with water. If changing from a solvent-based liquid to this water based flowable, thoroughly clean with methylated spirits then rinse with water. Failure to do so will result in clogging of equipment.

### **PRECAUTIONS**

DO NOT allow seed treated with this product to contaminate seed intended for human consumption. If possible, use a separate auger for treating seed. If using the same auger for treating seed and moving grain for human consumption, remove all AC Haven Flowable Seed Dressing residues from the auger to avoid contaminating untreated seed with seed dressing residues.

### **PROTECTION OF LIVESTOCK**

Seed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed. DO NOT allow seed treated with this product to contaminate seed intended for animal consumption.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT feed treated seed or otherwise expose to wild or domestic birds. Harmful to fish. DO NOT contaminate streams, rivers, or waterways with the chemical, used containers or bags which have held treated seed.

### **STORAGE AND DISPOSAL**

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product. DO NOT allow seed treated with this product to contaminate grain intended for human consumption. DO NOT use seed treated with this product for human consumption. Bags which have held treated seed are not to be used for any other purpose.

**For Non-Refillable Containers:** Store product in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse empty containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For Refillable Containers:** Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Axichem Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SAFETY DIRECTIONS**

May irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

## LABEL ELEMENTS

GHS Label Elements



**SIGNAL WORD** NOT APPLICABLE

### Hazard Statement(s)

<b>H373</b>	May cause damage to organs through prolonged or repeated exposure.
<b>H412</b>	Harmful to aquatic life with long lasting effects.
<b>H320</b>	Causes eye irritation.

### Precautionary statement(s) Prevention

<b>P260</b>	Do not breathe mist/vapours/spray.
<b>P273</b>	Avoid release to the environment.
<b>P264</b>	Wash all exposed external body areas thoroughly after handling.

### Precautionary statement(s) Response

<b>P305+P351+P338</b>	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
<b>P314</b>	Get medical advice/attention if you feel unwell.
<b>P337+P313</b>	If eye irritation persists: Get medical advice/attention.

### Precautionary statement(s) Storage

Not Applicable.

### Precautionary statement(s) Disposal

<b>P501</b>	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

## LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repackaging or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

**AxiChem**

**AxiChem**

**AxiChem**