

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 150 g/L INDOXACARB (s-isomer)

GROUP **22A** INSECTICIDE

For the control of various species of insect pests in Adzuki beans, Chickpeas, Cotton, Faba beans, Mungbeans and Soybeans as specified in the Directions for use table.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Returnable containers: (110L, 1000L)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist [or equivalent clothing] and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet which is available from the supplier.

## LIMIT OF WARRANTY AND LIABILITY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

APVMA Approval No: 87839/119532

Batch No:

Date of Manufacture:

**AxiChem**  
Pty Ltd

AXICHEM PTY LTD  
A.B.N 35 131 628 594

9 PALINGS COURT, NERANG QLD 4211  
PHONE: 07 5596 1736 www.axichem.com.au

**AxiChem**

## DIRECTIONS FOR USE

### Restraints

DO NOT apply if rain is expected within 2 hours of application, or if heavy dew is present on crops.

DO NOT apply when wind speed is less than 3 and greater than 20 kilometres per hour or during weather conditions when surface temperature inversions can develop.

DO NOT apply within 50 m (aerial application) or 20 m (ground application) when there are livestock, pasture or any land that is producing feed for livestock downwind from the application area.

**COTTON: DO NOT** apply more than three (3) applications per field in any one cotton growing season and no more than two (2) consecutive sprays per field per season. Applications must be a minimum of seven days apart.

**ADZUKI BEANS, CHICKPEAS, FABA BEANS, MUNGBEANS, SOYBEANS: DO NOT** apply more than one (1) application per field for the crops entire growth cycle.

### ENSURE YOU READ THE PROTECTION STATEMENTS BEFORE APPLYING THE PRODUCT.

CROP	INSECT	STATE	RATE
Cotton	Cotton bollworm ( <i>Helicoverpa armigera</i> ), Native budworm ( <i>H. punctigera</i> )	NSW, NT, Qld, WA only	650 mL/ha or 850 mL/ha
	Cotton bollworm ( <i>Helicoverpa armigera</i> ), Native budworm ( <i>H. punctigera</i> )		650 mL/ha + 2 L/ha Ovasyn* insecticide (or 200 g/L Amitraz EC formulation)
	Green mirid ( <i>Creontiades dilutus</i> )		650 mL/ha or 850 mL/ha or 300 or 400 mL/ha + salt (NaCl) at 5 g/L spray volume by ground (100 L/ha) or 10 g/L spray volume by air (30 L/ha)
Chickpeas, Faba beans	Cotton bollworm ( <i>H. armigera</i> ), Native budworm ( <i>H. punctigera</i> )	All States	300 mL/ha
Adzuki beans, Mungbeans, Soybeans	Cotton bollworm ( <i>H. armigera</i> ), Native budworm ( <i>H. punctigera</i> )	All States	400 mL/ha
	Mirid complex: Green mirid ( <i>C. dilutus</i> ), Brown mirid ( <i>C. pacificus</i> ), Crop mirid ( <i>Sidna kingbergii</i> ), Yellow mirid ( <i>Campylomma liebkechti</i> )		400 mL/ha + salt (NaCl) at 5 g/L spray volume by ground (100 L/ha) or 10 g/L spray volume by air (30 L/ha).
	Soybean looper ( <i>Thysanoplusia orichalcea</i> )		200 mL/ha
	Red shouldered leaf or Monolepta beetle ( <i>Monolepta australis</i> ) Soybeans only		200 mL/ha

## CRITICAL COMMENTS

Use the lower rate of G AC Jettison 150 EC Insecticide when:

- (1) *H. armigera* specific field levels are less than or equal to 60 % prior to treatment application AND
- (2) egg and larvae pressure ARE AT 5 – 10 brown eggs and 2 very small (first instar) or small larvae (second instar) per 10 cotton terminals AND
- (3) where preservation of beneficial insects is desirable.

Use the higher rate of AC Jettison 150 EC Insecticide when:

- (1) *H. armigera* specific field levels are greater than 60 % prior to treatment application AND
- (2) egg and larvae pressure ARE AT 5 – 15 brown eggs and 2 very small (first instar) or small larvae (second instar) per 10 cotton terminals AND
- (3) where preservation of beneficial insects is desirable.

Use AC Jettison 150 EC Insecticide + Ovasyn or 200 g/L Amitraz EC when:

- (1) egg and larvae pressure ARE AT 15 – 20 brown eggs and 2 very small (first instar) or small larvae (second instar) per 10 cotton terminals AND
- (2) where limited preservation of beneficial insects is required.

Target nymphs and/or adults when they reach the economic spray threshold. Use 650 or 850 mL/ha when controlling *Helicoverpa* spp. AND green mirids. Refer *Heliothis* recommendations. Use 300 or 400 mL/ha + salt when controlling green mirids ONLY. Use the higher rate on infestations exceeding economic spray threshold levels and/or large canopy crops.

**Under high populations suppression only may be observed.**

Note: AC Jettison 150 EC Insecticide has limited residual activity in controlling new infestations of mirids (either new hatchings of nymphs or influx of adults) post-spray.

Target brown eggs and hatchling (neonates or first instar) to small larvae (second instar) when they reach the economic spray threshold and before they become entrenched in flowers (particularly relevant to Faba beans) or pods.

Target brown eggs and hatchling (neonates or first instar) to small larvae (second instar) when they reach the economic spray threshold and before they become entrenched in flowers and pods.

Target nymphs and/or adults when they reach the economic spray threshold.

**Under high populations suppression only may be observed.** Please note: Gen AC Jettison 150 EC Insecticide has limited residual activity in controlling new infestations of mirids (either new hatchlings of nymphs or influx of adults) post spray.

Target hatchling (neonates or first instar) to small larvae (second instar) when they reach the economic spray threshold.

Target adult beetles when they reach the economic spray threshold. Ensure thorough spray coverage.

# NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

## WITHHOLDING PERIODS

### HARVEST

**COTTON: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.**

**ADZUKI BEANS, CHICKPEAS, FABA BEANS, MUNGBEANS, SOYBEANS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.**

### GRAZING

**COTTON: DO NOT ALLOW LIVESTOCK TO GRAZE CROPS, COTTON STUBBLE OR GIN TRASH TREATED WITH AC JETTISON 150 EC INSECTICIDE.**

**ADZUKI BEANS, CHICKPEAS, FABA BEANS, MUNGBEANS, SOYBEANS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.**

### LIVESTOCK DESTINED FOR EXPORT MARKETS

The label withholding period for grazing only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that the Export Slaughter Interval or the Export Grazing Intervals is observed before stock are sold or slaughtered.

### EXPORT SLAUGHTER INTERVAL (ESI): 28 DAYS

Livestock that have been grazing on or fed treated crops and/or over-sprayed should be placed on clean feed for 28 days (4 weeks) prior to export slaughter.

### GENERAL INSTRUCTIONS

AC Jettison 150 EC Insecticide is an emulsifiable concentrate formulation.

AC Jettison 150 EC Insecticide should be applied after careful field monitoring of pest populations of eggs and larvae to determine the need for application, the correct timing of the initial application and of any subsequent applications. For Cotton only, subsequent applications are dependent on economic thresholds, as well as the growth rate of new unprotected cotton terminals.

For *Helicoverpa* species, spray applications should be timed to coincide with egg hatching and before larvae are entrenched in protected feeding sites.

AC Jettison 150 EC Insecticide has been specifically designed for use in Integrated Pest Management schemes. The active ingredient, indoxacarb enters larvae primarily by ingestion of treated foliage, or through penetration of the insect cuticle. **After ingesting indoxacarb, the larvae cease feeding and die three to five days later.** AC Jettison 150 EC Insecticide does not give traditional larval "knockdown" control but controls nominated larvae species giving superior:

- square, flower and boll protection in cotton OR
- foliage, flower and pod protection in Chickpea, Faba beans, Mungbeans or Soybeans.

### PRODUCT USE

The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator must consider all these factors when making application decisions and determining off-target drift risks near the application. A spray drift minimisation strategy, should be employed at all times when applying this product.

APPLYING MEDIUM SIZE SPRAY DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT MINIMISE DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVOURABLE ENVIRONMENTAL CONDITIONS. Larger droplets may reduce the effects of evaporation.

### MIXING

Use only clean water. Half fill the spray tank with water and add the appropriate amount of AC Jettison 150 EC Insecticide directly to the spray tank, agitate and add Ovasyn or amitraz 200 g/L EC, (if applicable), then completely fill the tank. Mix thoroughly and continue mechanical or hydraulic agitation.

### Storage of spray mixture

Use the prepared spray immediately. If unforeseen conditions prevent immediate use of the AC Jettison 150 EC Insecticide spray mix, the mix may be stored up to 72 hours. Before use, thoroughly agitate the spray mix until fully resuspended. Mixtures of AC Jettison 150 EC Insecticide plus Ovasyn or Amitraz 200 g/L EC should not be stored.

## APPLICATION

Application equipment should be calibrated to apply at least sixty (60) droplets per cm<sup>2</sup> of target foliage. Droplets should be of MEDIUM spray droplet size.

### Ground application

Apply as a blanket spray or as a banded spray to all crops. Ensure thorough spray coverage on the foliage, using appropriate fan nozzles. Apply in a minimum spray volume of 100 L/ha and keep the boom low to avoid spray drift. A minimum spray pressure of 275 kPa (40 psi) should be used with fan nozzles applying insecticides.

Higher pressure reduces droplet size, **DOES NOT** improve canopy penetration AND MAY INCREASE DRIFT POTENTIAL. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE. For band spraying, increase the number of fan nozzles per crop row as the plant size increases.

### Aerial application

AC Jettison 150 EC Insecticide must only be applied with aircraft fitted with accurately calibrated equipment. Apply a minimum total spray volume of 30 L/ha with nozzles (e.g. Micronaire rotary atomisers, CP nozzles or conventional hydraulic nozzles) set to medium spray quality according to ASAE S572 definition for standard nozzles. A spray drift minimisation strategy, should be employed at all times when applying this product. DO NOT apply AC Jettison 150 EC Insecticide using Ultra Low Volume (ULV) methods.

### COMPATIBILITY

Label instructions for all products must be observed AC Jettison 150 EC Insecticide is compatible with Ovasyn insecticide, Amitraz 200 g/L EC formulations and Mepiquat 38 g/L.

AC Jettison 150 EC Insecticide is not compatible with ultra low volume (ULV) formulations or the following foliar fertilisers: Supa K 30, Zip and Triple 7. Since formulations may be changed and new ones introduced, it is recommended that users pre-mix a small quantity of the desired tankmix and observe possible adverse changes (settling out, flocculation etc).

### Salt

When the addition of salt is indicated in the Directions for Use table, Axichem Pty Limited recommends the use of salt from the following sources: Table or cooking salt, Pool salt, or salt approved for use in livestock feed e.g. Cheetham RAM (No. 2) Dried Fine Salt, Olsson's Kiln Dried Course Refined Salt.

### Spray Equipment Cleanout

Only apply product using clean, well maintained equipment. Immediately following application, thoroughly clean all spray equipment to reduce risk of deposits forming that might become difficult to remove.

Drain spray equipment into a disposal pit designed for this purpose. Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water. Fill the sprayer with clean water and household ammonia (one litre of 3% active for every 100 L of water). Flush hoses, boom, and nozzles. Turn off boom and top off the tank with clean water. Circulate through the spraying system for at least 15 minutes. Flush the hoses, boom and nozzles and drain the tank. Remove and clean nozzles, screens, and strainers in a bucket of fresh ammonia and water. Thoroughly rinse the sprayer, hoses, boom and nozzles with clean water several times. Clean all other associated contaminated application equipment.

### INSECTICIDE RESISTANCE WARNING

<b>GROUP</b>	<b>22A</b>	<b>INSECTICIDE</b>
--------------	------------	--------------------

For insecticide resistance management AC Jettison 150 EC Insecticide is a Group 22A insecticide. Some naturally occurring insect biotypes resistant to AC Jettison 150 EC Insecticide and other Group 22A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AC Jettison 150 EC Insecticide or other Group 22A insecticides are used repeatedly. The effectiveness of AC Jettison 150 EC Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Jettison 150 EC Insecticide to control resistant insects. Strategies to minimise the risk of insecticide resistance are available. To help prevent the development of resistance to AC Jettison 150 EC Insecticide observe the following instructions:

- Use AC Jettison 150 EC Insecticide in accordance with the current Insecticide Resistance Management (IRM) strategy for your region.
- Cultivate all cotton fields as soon as possible after picking to destroy overwintering pupae of *Helicoverpa armigera*.

For further information contact your local supplier, Axichem representative or local agricultural department agronomist.

## **PRECAUTION**

DO NOT use human flaggers/markers unless they are protected by engineering controls such as vehicles with enclosed cabs.

## **RE-ENTRY PERIOD**

DO NOT allow entry into treated areas until spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrists, a washable hat and chemical resistant gloves. Clothing must be laundered after each day's use.

## **PROTECTION OF LIVESTOCK**

**Dangerous to bees.** DO NOT apply when bees are actively foraging. Avoid direct application or drift of the spray mix onto beehives. After the spray has dried, bees can safely forage flowering crops.

## **AVOID SPRAY DRIFT ONTO ADJOINING PROPERTIES OR STOCK AREAS.**

Assess the treatment area before application to identify animal exposure risks. Avoid aerial application where possible. Observe the buffer zones for aerial and ground application. If unexpected conditions cause spray drift onto pasture or fodder crops that livestock may potentially graze or may be cut for livestock feed seek advice from Axichem Pty Limited.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Retain irrigation water and DO NOT allow the chemical to enter adjacent paddocks, crops or water supplies.

## **PROTECTION OF NON-TARGET BENEFICIAL INSECTS**

Beneficial insects contribute to control of secondary pest outbreaks. AC Jettison 150 EC Insecticide applications are unlikely to affect spiders and lacewings. Applications MAY temporarily reduce populations of predatory beetles, transverse ladybirds, ants and pirate bugs, but populations quickly recover.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby non-target plants/crops, cropping lands or pastures.

Refer to the Product Use section above and the cotton industry's Best Management Practice Manual to manage spray drift during application.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Returnable containers (110L, 1000L)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist [or equivalent clothing] and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

For further information, refer to the Safety Data Sheet which is available from the supplier.

**LABEL ELEMENTS**
**GHS Label Elements**

**SIGNAL WORD** Danger

**Hazard Statement(s)**

<b>H315</b>	Causes skin irritation.
<b>H317</b>	May cause an allergic skin reaction.
<b>H319</b>	Causes serious eye irritation.
<b>H372</b>	Causes damage to organs through prolonged or repeated exposure.
<b>H411</b>	Toxic to aquatic life with long lasting effects.
<b>H302</b>	Harmful if swallowed.

**Precautionary statement(s) Prevention**

<b>P260</b>	Do not breathe mist/vapours/spray.
<b>P280</b>	Wear protective gloves, protective clothing, eye protection and face protection.
<b>P264</b>	Wash all exposed external body areas thoroughly after handling.
<b>P270</b>	Do not eat, drink or smoke when using this product.
<b>P273</b>	Avoid release to the environment.
<b>P272</b>	Contaminated work clothing should not be allowed out of the workplace.

**Precautionary statement(s) Response**

<b>P302+P352</b>	IF ON SKIN: Wash with plenty of water.
<b>P305+P351+P338</b>	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
<b>P314</b>	Get medical advice/attention if you feel unwell.
<b>P333+P313</b>	If skin irritation or rash occurs: Get medical advice/attention.
<b>P337+P313</b>	If eye irritation persists: Get medical advice/attention.
<b>P362+P364</b>	Take off contaminated clothing and wash it before reuse.
<b>P391</b>	Collect spillage.
<b>P301+P312</b>	IF SWALLOWED: Call a POISON CENTER/doctor/physician/first aider if you feel unwell.
<b>P330</b>	Rinse mouth.

---

**Precautionary statement(s) Storage**

---

Not Applicable.

---

**Precautionary statement(s) Disposal**

---

**P501**

Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.

---

**LIMIT OF WARRANTY AND LIABILITY**

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.