

PEEL
HERE

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 250 g/L MCPA present as the ethyl hexyl ester, 25 g/L DIFLUFENICAN
SOLVENTS: 325 g/L LIQUID HYDROCARBONS, 150 g/L N-METHYL-2- PYRROLIDONE

GROUP F I HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in Directions for Use Table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

This leaflet is part of the label. READ THOROUGHLY BEFORE USE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site.

If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product.

For refillable containers (110L and 1000L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from Axichem Pty Ltd.

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No : 67032/54999

Batch Number:

Date of Manufacture:

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE
---	--

AxiChem
Pty Ltd

AXICHEM PTY LTD
A.B.N 35 131 628 594

9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

WEED LIST

WEED COMMON NAME	SCIENTIFIC NAME
Canola (Rape)	<i>Brassica napus</i>
Capeweed	<i>Arctotheca calendula</i>
Charlock	<i>Sinapis arvensis</i>
Chickweed	<i>Stellaria media</i>
Common Sowthistle (Milk Thistle)	<i>Sonchus oleraceus</i>
Corn Gromwell	<i>Buglossoides arvensis</i>
Cowvine	<i>Ipomoea lonchophylla</i>
Crassula	<i>Crassula</i> spp.
Deadnettle	<i>Lamium amplexicaule</i>
Dense-flower Fumitory	<i>Fumaria densiflora</i>
Dock	<i>Rumex</i> spp.
Doublegee (Spiny Emex)	<i>Emex australis</i>
Fat Hen	<i>Chenopodium album</i>
Fireweed	<i>Senecio</i> spp.
Fumitory	<i>Fumitory</i> spp.
Hedge Mustard	<i>Sisymbrium officinale</i>
Hexham Scent (King Island Melilot)	<i>Melilotus indicus</i>
Horehound	<i>Marrubium vulgare</i>
Hyssop Loosestrife	<i>Lythrum hyssopifolia</i>
Iceplant	<i>Mesembryanthemum</i> spp.
Indian Hedge Mustard	<i>Sisymbrium orientale</i>
London Rocket	<i>Sisymbrium irio</i>
Long Storksbill	<i>Erodium botrys</i>
Marshmallow	<i>Malva parviflora</i>
Mouse-eared Chickweed	<i>Cerastium glomeratum</i>

WEED COMMON NAME	SCIENTIFIC NAME
Night-scented Stock	<i>Matthiola longipetala</i>
Paterson's Curse	<i>Echium plantagineum</i>
Peppergrass	<i>Lepidium</i> spp.
Prickly Lettuce	<i>Lactuca serriola</i>
Purple Goosefoot	<i>Scleroblitum atriplicinum</i>
Rough Poppy	<i>Papaver hybridum</i>
Saffron Thistle	<i>Carthamus lanatus</i>
Scarlet Pimpernel	<i>Anagallis arvensis</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Skeleton Weed	<i>Chondrilla juncea</i>
Sorrel	<i>Rumex</i> spp.
Stemless Thistle	<i>Onopordum acaulon</i>
Toad Rush	<i>Juncus bufonius</i>
Tree Hogweed	<i>Polygonum patulum</i>
Turnip Weed	<i>Rapistrum rugosum</i>
Varigated Thistle	<i>Silybum marianum</i>
Vetch (Tares)	<i>Vicia sativa</i>
Volunteer Lupins	<i>Lupinus</i> spp.
Ward's Weed	<i>Carrichtera annua</i>
Wild Radish	<i>Raphanus raphanistrum</i>
Wild Turnip	<i>Brassica tournefortii</i>
Wireweed (Hogweed)	<i>Polygonum aviculare</i>

DIRECTIONS FOR USE:**RESTRAINTS:****DO NOT** apply to crop or weeds that are stressed due to dry or excessively moist conditions.**DO NOT** apply to crops under stress due to disease or insect damage.**DO NOT** apply to frost affected crops or if frosts are imminent.**DO NOT** apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE /HA
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover) PASTURE Newly Sown and Established Clover-based pasture, Clover for Hay and Seed Production	Wild Radish	Up to the 2 leaf stage and not more than 60mm in diameter	WA only	250mL
		Up to the 4 leaf stage and not more than 120mm in diameter	All States	500mL
		Up to the 6 leaf stage and not more than 150mm in diameter		750mL
		Up to the 8 leaf stage and not more than 180mm in diameter		1.0L
	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	Up to the 2 leaf stage and not more than 60mm in diameter	Qld only	500mL
		Up to the 4 leaf stage and not more than 120mm in diameter		750mL
		Up to the 6 leaf stage and not more than 150mm in diameter		1.0L
	London Rocket	Up to the 5 leaf stage and not more than 120mm in diameter	Qld only	750mL
	Ward's Weed		SA only	
	Capeweed	Up to the 2 leaf stage and not more than 60mm in diameter Up to the 4 leaf stage and not more than 120mm in diameter	All States	500mL
				1.0L
	Crassula	Up to the 2 leaf stage		500mL
				750mL

CRITICAL COMMENTS

CROP STAGE

Cereals

Up to 750mL (3 leaf to late tillering stage - Z13 to 30)

Over 750mL (5 leaf to late tillering stage - Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).

WA only:

DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).

Warning: AC SUNDOWNER may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to '**Crop Tolerance**' section of **General Instructions**).

Application is recommended prior to the eighth trifoliolate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils.

DO NOT apply to Annual Medics or Lucerne.

Warning: AC SUNDOWNER may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to '**Crop Tolerance**' section of **General Instructions**).

WEED STAGE

Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:

- high crop and weed density
- late season germinations
- abnormal weed growth (including early flowering), high rate of product (up to the maximum rate of application specified for that weed) may be required.

AC SUNDOWNER will not effectively control:

- regrowth of suppressed weeds;
- transplanted weeds;
- regrowth from rhizomes or roots;
- weeds growing under stress from previous herbicide applications.

GRAZING

Efficacy on larger weeds will be improved by grazing with normal levels of stock after 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions.

APPLICATION

Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased.

DIRECTIONS FOR USE: *continued*

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA		
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)	Prickly Lettuce	Up to the 2 leaf stage	All States	500mL		
		Up to the 4 leaf stage		750mL		
		Up to the 6 leaf stage		1.0L		
	Dense-flower Fumitory	Up to the 2 leaf stage		750mL		
				1.0L		
	Corn Gromwell, Saffron Thistle, Toad Rush	Up to the 2 leaf stage		NSW, Vic, SA only	1.0L	
					1.0L	
	Newly Sown and Established Clover-based pasture, Clover for Hay and Seed Production	Deadnettle		Up to the 2 leaf stage	Vic only	1.0L
		Sorrel		Up to the 4 leaf stage	All States	500mL
		Canola (Rape)		Up to the 6 leaf stage	Qld only	500mL
Purple Goosefoot		Cotyledon to 2 leaf	NSW only (West of Newell Hwy.) SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	350mL		
Turnip Weed Wild Turnip				500mL + 200mL terbutryn (500g/L)		
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Fumitory	2 - 6 leaf stage	All States	500mL + 200mL terbutryn (500g/L)		

SUPPRESSION OF THE FOLLOWING WEEDS

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)	Saffron Thistle	Up to the 6 leaf stage	All States	1.0L
	Chickweed, Fireweed, Hexham Scent (King Island Mellilot), Iceplant, Mouse-eared Chickweed, Night-scented Stock, Paterson's Curse, Peppergrass, Skeleton Weed, Long Storksbill, Volunteer Lupins.	Up to the 4 leaf stage		750mL
				1.0L
PASTURE Newly Sown and Established Clover-based pasture, Clover for Hay and Seed Production	Wireweed (hogweed)	Up to the 2 leaf stage		1.0L
Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee, (Spiny Emex), Fat Hen, Horehound, Hyssop Loosestrife, Marshmallow, Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Varigated Thistle, Vetch (Tares).				

CRITICAL COMMENTS

Continued.

WILD RADISH

AC SUNDOWNER will provide residual control of Wild Radish for up to 4 weeks after application.

Effective residual activity of this product may be reduced where:

- rates lower than 1.0L/ha are used;
- dry conditions prevail;
- poor coverage of the soil surface is achieved;
- crop is planted in non-wetting sand;
- soils contain a high content of organic matter.

Optimum results will be obtained if good soil moisture exists at and after application

CRITICAL COMMENTS

Refer to critical comments in the previous section.

DIRECTIONS FOR USE: *continued*

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Wild Radish	Up to the 4 leaf stage and not more than 120mm in diameter	All States	350mL plus 200mL L.V.E MCPA (500g/L)
		Up to the 6 leaf stage and not more than 150mm in diameter		500mL plus 200mL L.V.E MCPA (500g/L)
		Up to the 8 leaf stage and not more than 180mm in diameter*		500mL plus 400mL L.V.E MCPA (500g/L)

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD:

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

All Crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS:

RESISTANT WEEDS WARNING

GROUP F I HERBICIDE

AC SUNDOWNER is a member of the phenoxy and nicotinanalide groups of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management AC SUNDOWNER is both a Group F and a Group I herbicide. Some naturally occurring weed biotypes resistant to AC SUNDOWNER and other Group F and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by AC SUNDOWNER or other Group F or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of AC SUNDOWNER to control resistant weeds.

Tolerance

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Sub-clover is particularly sensitive.

Cereals

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

CRITICAL COMMENTS

Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above.

*Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180mm in diameter.

DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.

Crop Stage

AC SUNDOWNER 350mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30).

AC SUNDOWNER 500mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30).

AC SUNDOWNER 500mL + L.V.E. MCPA 400mL: Apply from 5 leaf stage to fully tillered (Zadoks Z15 to Z30).

Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post-sowing).

WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).

Warning: AC SUNDOWNER may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to 'Crop Tolerance' section of General Instructions).

Observe instructions also on L.V.E. MCPA product label.

Warning (Oats)

The tolerance of varieties Esk and Nile (the two main varieties grown in Tasmania) to AC SUNDOWNER has not been tested. Test a small area of crop before using AC SUNDOWNER over large areas. Consult your local Axichem Pty Ltd representative for advice on specific varieties.

Pasture

The tolerance of clover varieties to AC SUNDOWNER can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

Warning

AC SUNDOWNER may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500mL/ha and in areas of double spray. For this reason application is recommended prior to the 8th trifoliate leaf stage. However, at the lower rates (500mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to AC SUNDOWNER applied at 500mL/ha:

Arrow Leaf: Zulu

Berseem: Sacromonte

White: Haifa

Balansa: Paradana

Persian: Kyambro, Lupers, Maral

Subterranean Clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup. The effects of AC SUNDOWNER on clover seed yield have been tested on the following varieties. Under normal growing conditions they show levels of tolerance to AC SUNDOWNER applied at 500mL/ha.

Subterranean Clover: Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

Warning

Rose and Strawberry clover have shown increased sensitivity to AC SUNDOWNER. AC SUNDOWNER may affect the seed or yield of subterranean variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of AC SUNDOWNER. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties not listed should be tested before using AC SUNDOWNER over large areas. Consult your local Axichem Pty Ltd representative for advice on specific varieties.

Subsequent Crops

To reduce effect on susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

Mixing

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing AC SUNDOWNER should not be left to stand overnight.

Prolonged periods of exposure to cold temperature could result in settling out of the product in the mixture.

Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in AC SUNDOWNER and some other agricultural products. To reduce the risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

Application

Boom Sprayer

A minimum of 50L of water per hectare should be used, however, for optimum results water rates of 70-100L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

Selective Herbicide

Aircraft (NSW, Vic, SA only)

Apply in a minimum of 30L water per hectare. Effective control will only be achieved where good coverage of leaf surface is achieved.

Compatibility

The following products are physically compatible with AC SUNDOWNER as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	AC SUNDOWNER	Compatible Product
Wheat, triticale and cereal rye only	Up to 750mL/ha	Hoegrass (also barley), Puma Progress (wild oats only), Tristar Advance (1.5L rate only), Wildcat (wild oats only)
Cereals (including undersown)	All rates	Chlorpyrifos 500 EC Insecticide, Dimethoate 400 EC Insecticide, Endosulfan
Cereals (not undersown)	Up to 500mL/ha	AC Grimace 600 WG Herbicide, AC Dodge 750 Herbicide, MCPA (as 2-ethyl hexyl ester) 500 EC herbicide, AC Tulloch 750 Herbicide
	All rates	Bromoxynil (n-Octanoyl ester) 200 EC herbicide, Amine 625, AC Carnage, AC Frenzy Herbicide, dicamba (900g/kg) (up to 115 g only), Eclipse
Wheat, Barley, Triticale, and Cereal Rye only (not undersown)		AC Trotter 400WG Herbicide
Wheat only (not undersown)		AC Tussel Selective Herbicide
Clover	Up to 750mL/ha	AC Quiza 99.5EC Herbicide
Sub clover		Simazine Liquid, Simazine Liquid + Paraquat (200 g/L) mixture
	Up to 1.0L/ha	2,4-DB amine (500g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Hoegrass, Wildcat, Puma Progress or Tristar some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with dicamba dry a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties. Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. DO NOT add surfactant when mixing AC SUNDOWNER and AC Grimace 600 WG Herbicide.

Simazine: Refer to the simazine label for correct application rates, especially with regard to soil types. This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on the product: Sumi-Alpha Flex Insecticide, Alpha-Cypermethrin 100 EC insecticide, Karate and Bifenthrin 100 EC miticide or Bifenthrin 80 SC miticide.

Warning

DO NOT use crop oils with AC SUNDOWNER or AC SUNDOWNER mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of Axichem Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK:

Grazing Precaution

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site.

If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product.

For refillable containers (110L and 1000L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from Axichem Pty Ltd.

LABEL ELEMENTS

GHS Label Elements

**SIGNAL WORD** DANGER**Hazard Statement(s)**

H227	Combustible liquid.
H302	Harmful if swallowed.
H304	May be fatal if swallowed and enters airways.
H312	Harmful in contact with skin.
H315	Causes skin irritation.
H319	Causes serious eye irritation.
H332	Harmful if inhaled.
H335	May cause respiratory irritation
H336	May cause drowsiness or dizziness.
H351	Suspected of causing cancer.
H360	May damage fertility or the unborn child.
H410	Very toxic to aquatic life with long lasting effects.

Precautionary statement(s) Prevention

P201	Obtain special instructions before use.
P210	Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
P271	Use only outdoor or in a well-ventilated area.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P281	Use personal protective equipment as required.
P261	Avoid breathing mist/vapours/spray.
P264	Wash all exposed external body areas thoroughly after handling.
P270	Do not eat, drink or smoke when using this product.
P273	Avoid release to the environment.

Precautionary statement(s) Response

P301+P310	IF SWALLOWED: Immediate call a POISON CENTRE/doctor/physician/first aider.
P331	Do NOT induce vomiting. If more than 15 mins from Doctor, INDUCE VOMITING (if conscious).
P308+P313	If exposed or concerned: Get medical advice/attention.
P370+P378	In case of fire: Use alcohol resistant foam or normal protein foam to extinguish.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+P313	If eye irritation persists: Get medical advice/attention.
P391	Collect spillage.
P301+P312	IF SWALLOWED: Call a POISON CENTRE or Doctor/Physician if you feel unwell.
P302+P352	IF ON SKIN: Wash with plenty and water.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P330	Rinse mouth.
P332+P313	If skin irritation occurs: Get medical advice/attention.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

P405	Store locked up.
P403+P233	Store in a well-ventilated place. Keep container tightly closed.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

AxiChem

AxiChem