


PEEL
HERE

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 240g/L OXYFLUORFEN
SOLVENTS: 606g/L LIQUID HYDROCARBON
108g/L N-METHYL-2- PYRROLIDONE

GROUP G HERBICIDE

For the selective control of certain broadleaf weeds and some grasses as per the directions for use table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For REFILLABLE containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the Container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hand after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia: 131126, New Zealand 0800 774 766. If swallowed, DO NOT induce vomiting. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS).

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repackaging or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval Number: 68742/128093

Batch No.:

Date of Manufacture:

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY
CALL 1800 039 008
ALL HOURS AUSTRALIA WIDE

AxiChem
Pty Ltd

AXICHEM PTY LTD
A.B.N 35 131 628 594

9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

DIRECTIONS FOR USE:

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tree fruit, Nuts and Vines not less than 3 years old DORMANT APPLICATION Almonds, Apples, Apricots, Grapevine, Macadamias, Peaches, Plums, Pears, Pecans, Walnuts	Refer to Weeds Controlled list	Weed free soil	3 or 4L/ha	DO NOT apply once bud swell has occurred. Apply to freshly cultivated weed free soil. Use the higher rate when longer residual control is required (up to 4 months). Where grass weeds are expected to be the major problem, or when control of a wider weed spectrum is required, mix the lower rate with 4-5 L Oryzalin (500g/L) per treated hectare. Read and follow all label directions.
		Emerged weeds present		DO NOT apply once bud swell has occurred. Use the higher rate when longer residual control is required (up to 4 months). When light infestations of weeds are present, apply as a tank mix with Glyphosate (450g/L) or paraquat (250g/L) to give both knockdown and residual control. Read and follow all label directions. Where weed growth is large and dense, weeds must be eliminated prior to application of AC Oxxel 240 EC Herbicide, using glyphosate (450g/L) or mechanical means.
	Annual Weeds Refer to Glyphosate 450g/L label	Actively growing emerged weeds	75 mL/ha plus Glyphosate 450g/L at recommended label rates	Addition of AC Oxxel 240 EC Herbicide to Glyphosate (450g/L) will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with Glyphosate 450g/L applied alone) and give control of annual nettles <i>Urtica spp.</i> , and Marshmallow, <i>Malva parviflora</i> . For rates refer to the appropriate label. Read and follow all label directions. DO NOT apply the tank mix of AC Oxxel 240 EC Herbicide and Glyphosate (450g/L) near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
<p>Tree fruit, Nuts and Vines not less than 3 years old</p> <p>DORMANT APPLICATION</p> <p>Almonds, Apples, Apricots, Grapevine, Macadamias, Peaches, Plums, Pears, Pecans, Walnuts</p>	<p>Most annual grasses and broadleaf weeds. Refer to Weeds controlled list and Paraquat/Diquat or Paraquat only labels</p>	<p>Emerged weeds present</p>	<p>250mL plus Paraquat/Diquat or Paraquat only at recommended label rates</p> <p>OR</p> <p>25 to 30mL/100L plus Paraquat/Diquat or Paraquat only at recommended label rates</p>	<p>The addition of AC Oxxel 240 EC Herbicide to paraquat/diquat or paraquat only will improve control of small flowered mallow, evening primrose and other weeds sensitive to AC Oxxel 240 EC Herbicide.</p> <p>Apply as a tank mix and refer to the appropriate label and follow all label directions.</p>
<p>Duboisia</p>	<p>Refer to weeds controlled list</p>	<p>Weed free soil</p>	<p>4 or 8L/ha</p>	<p>Apply after transplantation of young seedlings or after harvest of mature plants. Use higher rate where longer residual control (up to 6 months) is required. "Over-the-top" application will be tolerated.</p>
		<p>Weeds at the 4 to 6 true leaf stage</p>		<p>Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Established larger weeds must be eliminated by mechanical means prior to the application. Addition of a non-ionic surfactant (1000g/L) at recommended rates will improve activity.</p>
<p>Tropical and Subtropical fruit (inedible peel)</p> <p>Custard Apple, Kiwi Fruit, Longan, Lychees, Mango, Passionfruit, Pawpaw, Rambutan</p>	<p>Refer to Weeds Controlled list</p>	<p>Weed free soil</p>	<p>4 L/ha</p>	<p>Best results are obtained when applied to moist soil free of weeds.</p>
		<p>Emerged weeds present</p>		<p>If weeds are present AC Oxxel 240 EC Herbicide should be applied as a tank mix with the recommended rates of Glyphosate 450g/L, paraquat/diquat or paraquat.</p>

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Brassica Crops, Broccoli, Cabbages, Cauliflower	Refer to weeds controlled list	Weed free soil	1.5 to 2L/ha	Apply AC Oxxel 240 EC Herbicide to prepared ground 4 to 7 days prior to transplanting. If soil is dry irrigation or rainfall is required prior to transplanting for activation of AC Oxxel 240 EC Herbicide. Utilise transplanting techniques which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicide activity. Use the higher rate in situations where weed pressure is known to be heavy.
Onion (seeded)	Blackberry Nightshade (<i>Solanum nigrum</i>)	Hook leaf stage	50 to 100mL/ha	Do not exceed 100 mL/ha at the hook leaf stage as excessive crop damage may occur.
	Common Cotula (<i>Cotula australis</i>), Crowsfoot / Storksbill (<i>Erodium spp.</i>), Deadnettle (<i>Lamium amplexicaule</i>), Docks (<i>Rumex spp.</i>), Fumitory (Pink and white flowered) (<i>Fumaria spp.</i>), Groundsel (<i>Senecio vulgaris</i>), Hogweed / Wireweed (<i>Polygonum aviculare</i>), Milk Thistle (<i>Sonchus oleraceus</i>), Plantain (<i>Plantago spp.</i>), Potato weed (<i>Galinsoga parviflora</i>), Sorrel (<i>Rumex acetosella</i>), Volunteer potato (<i>Solanum tuberosum</i>)	1 ½ to 2 ½ leaf stage	350 - 500mL/ha	Best results are obtained when the hook leaf stage treatment is followed by a further application of AC Oxxel 240 EC Herbicide when onions are at the 1 ½ to 2 ½ leaf stage. Apply AC Oxxel 240 EC Herbicide in at least 100L/ha water at pressures below 150kPa. DO NOT exceed 150kPa because excessive crop damage could result. Multiple treatments may be applied provided the total dose does not exceed 2L/ha in one season. For control of other weeds, use in a program with other registered herbicides. NOTE: Can cause flecking, twisting or stunting of onion plants although such symptoms are normally short-lived. DO NOT apply in conditions of very high humidity. DO NOT apply to weeds or crops which are stressed by prevailing weather conditions, disease or mechanical damage. DO NOT use wetting agents or other adjuvants.

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tobacco	Refer to weeds controlled list	Weed free soil	4L	Use to control weeds along spray lines only. DO NOT apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. AC Oxxel 240 EC Herbicide should be applied to a moist soil. Where very small weeds (2 to 3 leaf) emerge prior to spraying, the addition of a wetting agent to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2 L/ha paraquat 200g/L is required. Avoid spray drift.
Pyrethrum – as bare rooted transplants or seedlings	Refer to weeds controlled list (except Chickweed)	Pre-Plant incorporated into weed free soil worked to a fine tilth	4 to 6L/ha	Apply prior to final soil preparation. The preferred implements for final soil preparation would be either a multiple tyne cultivator or rotary harrows. Use the 6L rate for heavy black clay soils only (as found in the Derwent and Coal River Valleys). AC Oxxel 240 EC Herbicide will not provide consistent control of chickweed.
Pyrethrum - greater than 4 leaf	Blackberry nightshade (<i>Solanum nigrum</i>), Cleavers (<i>Galium aparine</i>), Field Bindweed (<i>Convolvulus vulgaris</i>), Fumitory (<i>Fumaria spp.</i>), Groundsel (<i>Senecio vulgaris</i>), Hogweed (<i>Polygonum aviculare</i>), Sorrel (<i>Rumex acetosella</i>), Volunteer potato (<i>Solanum tuberosum</i>)	Emerged weeds present	100 – 150mL/ha	Apply when pyrethrum growth stage is greater than 4 true leaf.
Pyrethrum - established crops, greater than 1 year old	As above plus Sow Thistle (<i>Sonchus spp.</i>), Spear Thistle (<i>Cirsium vulgare</i>)		200mL to 4L/ha	Apply when pyrethrum is greater than 100 mm rosettes. Apply rates of greater than 1.0 L/ha ONLY between 1st of February and 31st of March. DO NOT apply later than 25 weeks before harvest.

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Prior to sowing winter cereals, cotton	Annual Weeds Refer to Weeds Controlled list and Glyphosate 450g/L or 360g/L labels	Actively growing emergent weeds	75mL/ha plus glyphosate 450g/L or glyphosate 360g/L at recommended label rates	Addition of AC Oxxel 240 EC Herbicide to Glyphosate 450 or Glyphosate 360 will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with Glyphosate 450g/L or 360g/L applied alone) and give control of annual nettles (<i>Urtica spp.</i>), and Marshmallow (<i>Malva parviflora</i>).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

GRAZING: DO NOT GRAZE TREATED WEEDS

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

Weeds Controlled By AC Oxxel 240 EC Herbicide include the Following:

WEEDS	SEEDLINGS (less than 4 true leaf)
Amsinckia <i>Amsinckia</i> spp.	Redshank <i>Amaranthus cruentus</i>
Barley grass <i>Hordeum leporinum</i>	Ryegrass <i>Lolium</i> spp.
Barnyard grass <i>Echinochloa</i> spp.	Sesbania pea <i>Sesbania cannabina</i>
Blackberry nightshade <i>Solanum nigrum</i>	Shepherd's purse <i>Capsella bursa-pastoris</i>
Bladder ketmia <i>Hibiscus trionum</i>	Smallflower mallow <i>Malva parviflora</i>
Burrgrass <i>Cenchrus australis</i>	Soursob <i>Oxalis pes-caprae</i>
Caltrop <i>Tribulus terrestris</i>	Sowthistle <i>Sonchus oleraceus</i>
Capeweed <i>Arctotheca calendula</i>	Starburr <i>Acanthospermum hispidum</i>
Chickweed <i>Stellaria media</i>	Stinkgrass <i>Eragrostis cilianensis</i>
Crowsfoot grass <i>Eleusine indica</i>	Summer grass <i>Digitaria</i> spp.
Deadnettle <i>Lamium amplexicaule</i>	Thornapple <i>Datura stramonium</i>
Fat hen <i>Chenopodium album</i>	White eye <i>Richardia brasiliensis</i>
Giant pigweed <i>Trianthema portulacastrum</i>	Wild mustard <i>Sisymbrium</i> spp.
Liverseed grass <i>Urochloa panicoides</i>	Wild radish <i>Raphanus raphanistrum</i>
Lovegrass <i>Eragrostis</i> spp.	Wireweed <i>Polygonum aviculare</i>
Pigeon grass <i>Setaria</i> spp.	
Pigweed <i>Portulaca oleracea</i>	
Prickly lettuce <i>Lactuca</i> spp.	
Red natal grass <i>Rhynchelytrum repens</i>	

GENERAL INSTRUCTIONS

TIMING

For optimum residual weed control, AC Oxxel 240 EC Herbicide should be applied to the soil surface prior to weed emergence after all other agricultural operations, such as mechanical cultivation and reshaping of irrigation furrows, have been completed. The area should be left undisturbed during the period for which weed control is desired. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

MIXING

When using AC Oxxel 240 EC Herbicide alone, fill the spray tank, at least one-third full with clean water, add the recommended amount of oxyfluorfen while the pump and agitator are running, then complete filling the spray tank. A non-ionic surfactant, if required by label directions, should be added near the end of the filling process to minimise foaming. When tank mixing with Oryzalin, add to one-third filled tank, then the AC Oxxel 240 EC Herbicide during the filling operation. Maintain agitation during mixing and until spraying is completed. When tank mixing with Glyphosate (450 g/L or 360 g/L), paraquat/diquat or paraquat formulations, add these after AC Oxxel 240 EC Herbicide during the filling operation. When using AC Oxxel 240 EC Herbicide in mixtures always refer to the appropriate label and read and follow all label directions.

APPLICATION

Spray equipment should be calibrated before use. AC Oxxel 240 EC Herbicide should be applied uniformly as a directed treatment to the base of tree and vine crops using flat fan or off-centre nozzle. Complete coverage of seedling weeds is required for maximum knockdown effect. A water volume of 250 to 500 Litres per hectare is recommended for treatments of bare soil. A spray volume of 100 to 1350 Litres per hectare is recommended where seedling weeds (4 to 6 leaf) are present. Ensure both weed foliage and the soil surface are sprayed. Use the higher volumes where weed density is high. Tank mixtures of 75mL/ha of AC Oxxel 240 EC Herbicide with glyphosate (450g/L or 360g/L) should be applied in 30 to 200 Litres spray volume per hectare. For maximum residual control, AC Oxxel 240 EC Herbicide should NOT be incorporated or disturbed after application.

CROP SAFETY

AC Oxxel 240 EC Herbicide may be applied as directed around dormant peach, plum, apricot, almond, apple and pear trees and grape vines of all ages when applied at rates of less than 1.0 L/ha. When applied at 3.0 L/ha and above, the trees and grape vines should be at least 3 years of age. Do NOT apply AC Oxxel 240 EC Herbicide once bud swell has occurred when using rates greater than 1.0 L/ha. Duboisia seedlings and mature plants will tolerate "over-the-top" applications of AC Oxxel 240 EC Herbicide. When using the 75 mL/ha rate, AC Oxxel 240 EC Herbicide may be applied as directed prior to sowing winter cereals.

COMPATIBILITY

AC Oxxel 240 EC Herbicide is compatible with Glyphosate 450g/L and 360g/L, paraquat, oryzalin and diquat.

RESISTANT WEEDS WARNING

GROUP	G	HERBICIDE
--------------	----------	------------------

AC Oxxel 240 EC Herbicide is a member of the diphenyl ether group of herbicides. AC Oxxel 240 EC Herbicide has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management AC Oxxel 240 EC Herbicide is a Group G herbicide. Some naturally occurring weed biotypes resistant to AC Oxxel 240 EC Herbicide and other Group G herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by

AC Oxxel 240 EC Herbicide or other Group G herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Oxxel 240 EC Herbicide to control resistant weeds.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear chemical resistant footwear.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby water bodies, susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is highly toxic to wildlife and fish. DO NOT contaminate lakes, ponds, streams, rivers or waterways with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For REFILLABLE containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the Container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hand after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia: 131126, New Zealand 0800 774 766. If swallowed, DO NOT induce vomiting. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS) which is available from the supplier.

GHS Label Elements



SIGNAL WORD	DANGER
--------------------	---------------

Hazard Statement(s)	
----------------------------	--

H304	May be fatal if swallowed and enters airways.
H315	Causes skin irritation.
H319	Causes serious eye irritation.
H336	May cause drowsiness or dizziness.
H351	Suspected of causing cancer.
H360	May damage fertility or the unborn child.
H373	May cause damage to organs through prolonged or repeated exposure.
H410	Very toxic to aquatic life with long lasting effects.

Precautionary statement(s) Prevention	
--	--

P201	Obtain special instructions before use.
P260	Do not breathe mist/vapours/spray.
P271	Use only outdoors or in a well-ventilated area.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P273	Avoid release to the environment.
P264	Wash all exposed external body areas thoroughly after handling.

Precautionary statement(s) Response	
--	--

P301+P310	IF SWALLOWED: Immediate call a POISON CENTRE/doctor/physician/first aider.
P331	Do NOT induce vomiting.
P308+P313	IF exposed or concerned: Get medical advice/attention.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P312	Call a POISON CENTER/doctor/physician/first aider/if you feel unwell.
P337+P313	If eye irritation persists: Get medical advice/attention.
P391	Collect spillage.

P302+P352	IF ON SKIN: Wash with plenty of water.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P332+P313	If skin irritation occurs: Get medical advice/attention.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

P405	Store locked up.
P403+P233	Store in a well-ventilated place. Keep container tightly closed.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

AxiChem

AxiChem