


CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 700 g/kg IMAZAMOX

GROUP B HERBICIDE

For the post-emergent control of certain annual grass and broadleaved weeds in field peas, legume-based pastures, Lucerne, peanuts and soybeans, as specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. When opening the container and preparing spray, wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet which is available from the supplier.

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repackaging or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 88755/122191

Batch No:

Date of Manufacture:

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

**FOR SPECIALIST ADVICE IN
AN EMERGENCY ONLY
CALL 1800 039 008
ALL HOURS AUSTRALIA WIDE**

Axi Chem
Pty Ltd

AXICHEM PTY LTD

A.B.N 35 131 628 594

9 PALINGS COURT, NERANG QLD 4211

PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to crops or weeds stressed by factors such as root or foliar diseases, water logging, nutrient deficiencies or extremes of temperature and moisture.

DO NOT apply to crops stressed by any previous herbicide treatments.

CROP	WEEDS CONTROLLED	RATE/ha
Field peas Post-emergence (See CROP SAFETY re varieties)	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus</i> spp.), Deadnettle (<i>Lamium amplexicaule</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Storksbill (<i>Erodium</i> spp.), Turnip weed (<i>Rapistrum rugosum</i>), Volunteer barley (<i>Hordeum vulgare</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer lupins (<i>Lupinus</i> spp.), Volunteer triticale (<i>Triticosecale</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Wild oats (<i>Avena</i> spp.), Wild turnip (<i>Brassica tournefortii</i>), *Doublegee (<i>Emex australis</i>), *Shepherds purse (<i>Capsella bursa-pastoris</i>), *Threehorn bedstraw (<i>Galium tricornutum</i>), *Wild radish (<i>Raphanus raphanistrum</i>), Wireweed (<i>Polygonum aviculare</i>)	45 g plus BS 1000 or equivalent, at 200 mL/100 L water

WHP	CRITICAL COMMENTS
6 weeks (grazing)	<p>DO NOT apply to field peas beyond the 4 node stage.</p> <p>DO NOT use adjuvants other than BS 1000 or equivalent.</p> <p>Apply to actively growing broadleaved weeds in the cotyledon to 3 leaf stage.</p> <p>Apply to grass weeds up to the 2 tiller stage.</p> <p>Good crop growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded.</p> <p>*Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds.</p> <p>Leaf yellowing and height suppression may occur, especially under adverse growing conditions such as moisture stress. Flowering may be delayed.</p> <p>IMPORTANT: Refer to CROP SAFETY section of label regarding use on field peas.</p> <p>Refer to FOLLOW CROPS section of this label regarding follow crops.</p>

CROP	WEEDS CONTROLLED	RATE/ha
Legume based pastures Post-emergence: Clovers, Lucerne, medics, serradellas only. See CROP SAFETY re varieties	As for field peas above.	45 g plus Hasten or Kwickin or equivalent at 500 mL/100 L water
	* Silver grass (<i>Vulpia</i> spp.) plus above weeds	50 g plus Hasten or Kwickin or equivalent at 500 mL/100 L water plus ammonium sulphate adjuvant at 2 L /100 L water
Lucerne Post-emergence. See CROP SAFETY re varieties	As for post emergence use in field peas (winter weeds) and soybeans (summer weeds).	45 g or 50 g plus BS 1000 or equivalent at 200 mL/100 L water plus ammonium sulphate adjuvant at 2 L /100 L water

WHP	CRITICAL COMMENTS
7 days (grazing)	<p><u>Seedling</u> Apply to legumes after full emergence of the third trifoliate leaf but well before commencement of flowering.</p> <p><u>Established</u> Apply before commencement of flowering. Apply to actively growing broadleaved weeds in the cotyledon to 3 leaf stage. Apply to grass weeds up to the 2 tiller stage. Good pasture growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Where pasture stand is poor, weed control may be inadequate due to lack of competition. Ensure sufficient spray coverage of weeds particularly in dense pastures. *Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Very high populations may not be efficiently controlled. Good spray coverage is essential. Ensure that crop agronomy is correct, to provide competition to such weeds. Refer to CROP SAFETY section of this label regarding varieties of clovers and medics and other pasture species. Note: Use of the 50 g /ha rate plus Hasten or Kwicken may severely damage medics. Refer to FOLLOW CROPS section of this label regarding follow crops.</p>
	<p>Apply to actively growing weeds at growth stages as for field peas (winter weeds) and soybeans (summer weeds). DO NOT use the Hasten or Kwicken mix on seedling Lucerne. Use the 50 g rate for summer weeds.</p> <p>Crop Stage <u>Seedling</u> Apply when crop is at 2 trifoliate leaf stage and before commencement of flowering. <u>Established</u> Apply as above following cutting or grazing. Refer to CROP SAFETY section of this label regarding varieties of Lucerne.</p>

CROP	WEEDS CONTROLLED	RATE/ha
Post-emergence Peanuts, Soybeans	Amaranth (<i>Amaranthus</i> spp.), Barley grass (<i>Hordeum leporinum</i>), Barnyard grass (<i>Echinochloa crus-galli</i>), Bell vine (<i>Ipomoea plebeian</i>), Brome grass (<i>Bromus</i> spp.), Deadnettle (<i>Lamium amplexicaule</i>), Fathen (<i>Chenopodium album</i>), Fierce thornapple (<i>Datura ferox</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Liverseed grass (<i>Urochloa panicoides</i>), Turnip weed (<i>Rapistrum rugosum</i>), Volunteer barley (<i>Hordeum vulgare</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer lupins (<i>Lupinus</i> spp.), Volunteer triticale (<i>Triticosecale</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Wild gooseberry (<i>Phsalis minima</i>), Wild oats (<i>Avena</i> spp.), Wild turnip (<i>Brassica tournefortii</i>), *Anoda weed (<i>Anoda cristata</i>), *Awneless barnyard grass (<i>Echinochloa colona</i>), *Blackberry nightshade (<i>Solanum nigrum</i>), *Caltrop (<i>Tribulus terrestris</i>), *Chickweed (<i>Stellaria media</i>), *Crabgrass (<i>Digitaria ciliaris</i>), *Doublegee (<i>Emex australis</i>), Noogoora burr (<i>Xanthium pungens</i>), *Shepherds purse (<i>Capsella bursa-pastoris</i>), *Threehorn bedstraw (<i>Galium tricornutum</i>), *Wild radish (<i>Raphanus raphanistrum</i>), Wireweed (<i>Polygonum aviculare</i>)	50 g plus EITHER BS 1000 or equivalent at 200 mL/100 L water plus ammonium sulphate adjuvant at 2 L /100 L water OR Hasten or Kwikin or equivalent at 500 mL or 1 L/100 L water

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WHP	CRITICAL COMMENTS
4 weeks (grazing)	<p>DO NOT apply to crops beyond the 6 leaf stage.</p> <p>Apply to actively growing weeds in the cotyledon to 2 leaf stage. Apply to grass weeds up to the 2 tiller stage.</p> <p>If using the Hasten or Kwickin mix, use the 1 L / 100 L rate where grass weeds predominate.</p> <p>*Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds.</p> <p>Refer to FOLLOW CROPS section of this label regarding follow crops.</p>

WITHHOLDING PERIODS

GRAZING

LUCERNE, PASTURES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

PEANUTS, SOYBEANS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION

FIELD PEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

HARVEST

FIELD PEAS, PEANUTS, SOYBEANS: NOT REQUIRED WHEN USED AS DIRECTED

RESISTANT WEED WARNING

GROUP	B	HERBICIDE
-------	---	-----------

AC REPO 700 WG HERBICIDE is a member of the imidazolinone group of herbicides. The product has the ALS mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50 m of wetlands or waterways.

Crop Safety: This product may cause shortening of plant internodes and may in some circumstances lead to transient crop yellowing but plants will soon recover. This effect is more pronounced under poor growth conditions including conditions of prolonged moisture stress.

Field peas

DO NOT apply to field peas beyond the 4 node stage.

DO NOT apply if frost is forecast.

DO NOT use AC REPO 700 WG HERBICIDE after simazine (unregistered in field peas) and do NOT exceed 250g ai/ha diuron if use of AC REPO 700 WG HERBICIDE is planned.

DO NOT roll field peas after germination if use of AC REPO 700 WG HERBICIDE is planned.

Before using this product on field pea varieties other than Bohatyr, Bonzer, Dun, Glenroy, Laura, Magnet, Mukta, Santi and Wirrega, contact your agronomist for up to date information. DO NOT use on Alma, Excell and Parafield varieties.

Pasture legumes and perennial grasses

Safety to medic varieties may be lower than to clover varieties. Safety to seedling pastures less than one year old may be lower than to older established pastures. Transient biomass reduction and leaf yellowing may occur to pasture legumes and perennial grasses, with good recovery occurring under favourable seasonal conditions following application. For up to date information on all pasture clover and medic varieties and other temperate pasture species, contact your local agronomist before use.

Lucerne

Before using this product on lucerne varieties other than Hunterfield, Sceptre, Sequel, Siriver and Trifecta, contact your local agronomist for up to date information

DO NOT USE THE HASTEN OR KWICKIN (or equivalent) MIX ON SEEDLING LUCERNE.

Replanting: If replanting is necessary in a field previously treated with AC REPO 700 WG HERBICIDE, the field may be replanted to field peas (above listed varieties only), soybeans or peanuts. DO NOT apply a second treatment of AC REPO 700 WG HERBICIDE.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic flora. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. When opening the container and preparing spray, wear face shield or goggles. If product in eyes, wash it out immediately with water, Wash hands after use. After each day's use, wash face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet which is available from the supplier.

LABEL ELEMENTS

GHS Label Elements	 
--------------------	---

SIGNAL WORD	WARNING
Hazard Statement(s)	
H315	Causes skin irritation.
H319	Causes serious eye irritation.
H335	May cause respiratory irritation.
H410	Very toxic to aquatic life with long lasting effects.

Precautionary statement(s) Prevention

P271	Use only outdoor or in a well-ventilated area.
P261	Avoid breathing dust/fumes.
P273	Avoid release to the environment.
P280	Wear protective glove, protective clothing, eye protection, and face protection.
P264	Wash all exposed external body areas thoroughly after handling.

Precautionary statement(s) Response

P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P312	Call a POISON CENTRE/doctor/physician/first aider/ if you feel unwell.
P337+P313	If eye irritation persists: Get medical advice/attention.
P391	Collect spillage.
P302+P352	IF ON SKIN: Wash with plenty of water.

P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P332+P313	If skin irritation occurs: Get medical advice/attention.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

P405	Store locked up.
P403+P233	Store in a well-ventilated place. Keep container tightly closed.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any state and Federal Legislation or law of a Territory. Not for repacking or reformulations. No license under any non-Australian patent is granted or implied by purchase of this container.

AxiChem