

PEEL
HERE

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 18 g/L ABAMECTIN
Solvents: 265 g/L N-METHYL-2-PYRROLIDONE
58 g/L LIQUID HYDROCARBON

GROUP 6 INSECTICIDE

For the control of certain mites on apples, capsicums, citrus, cotton, hops, ornamentals, pears, tomatoes and strawberries and Native Budworm on cotton.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT
This leaflet is part of the label. READ THOROUGHLY BEFORE USE

STORAGE AND DISPOSAL

Store below 30°C (Room temperature). Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture empty containers and deliver for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. For space applications, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles and a half face respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles, respirator (and if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from AxiChem Pty Ltd.

LIMIT OF WARRANTY AND LIABILITY

AxiChem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of AxiChem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excluding conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 66904/145363

Batch Number:

Date of Manufacture:

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE
--	---

AxiChem
Pty Ltd

AXICHEM PTY LTD
A.B.N 35 131 628 594
9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

DIRECTIONS FOR USE:**RESTRAINTS:**

DO NOT spray if rain seems likely before spray has dried on the leaves. Failure to observe this restraint may result in reduced efficacy.

SITUATION	PEST	RATE/ha	WHP (days)
Apples, Pears	Two Spotted Mite (<i>Tetranychus urticae</i>) European Red Mite (<i>Panonychus ulm</i>)	750 mL/ha plus 5L/ha summer oil See general instructions for mixing rates	14
Capsicums	Two Spotted Mite (<i>Tetranychus urticae</i>)	300 mL/ha	3
Citrus	Broad Mite (<i>Polyhagotarsonemus latus</i>) Brown Citrus Rust Mite (<i>Tegolophus australis</i>) Citrus Rust Mite (<i>Phyllocoptera oleivora</i>)	Dilute spray: 15mL/100mL or 25mL/100mL plus 250mL summer oil /100L Concentrate spray: Refer to the Application Section - Citrus	7

CRITICAL COMMENTS

For best results, apply the product using dilute application (spraying to point of run-off). If applying with concentrate applications efficacy may be comprised. DO NOT apply at more than two times concentrate (ie half the dilute water volume). Apply the same amount of AC WHISTLER Miticide/Insecticide to the target crop whether by spraying dilute or concentrate spray methods. DO NOT apply this product before or after applications of Delan* or Captan*.

In apples apply this product from 2 to 6 weeks after petal fall if monitoring shows high numbers of overwintering European Red Mite Eggs are present or if mites are a problem early in the season.

In pears, timing is not as critical and the application should be made as soon as after mite numbers have reached a threshold for your area.

Maximum mite control is usually reached at 7 days. Moderate to high mite populations will be controlled but if there are no predatory mites present re-treatment with another miticide (from another chemical group) may be necessary.

Integrated pest control: The effects of abamectin on parasitic wasps and other beneficial insects in Australian orchards are not known. Studies have shown that after application of abamectin predatory mite populations may not increase for a number of weeks, due to the lack of suitable pest mite prey. Predatory mite numbers will increase with any increase in pest mite numbers allowing the continuation of biological mite control. DO NOT use in IPM programmes unless the pest mite threshold has been reached and predators are unlikely to achieve effective control.

Thorough coverage and penetration into the plant canopy is essential. Preferably apply before the build-up of plant numbers. Re-apply as pest numbers indicate. Alternate with other chemical groups. Allow at least 28 days between applications with AC WHISTLER Miticide/Insecticide and do not make more than 2 AC WHISTLER Miticide/Insecticide sprays per crop. Refer to notes on resistance in the General Instructions section of this label.

Apply as a dilute spray in 3000 to 6000 L water/ha as pest pressure indicates.

Apply by dilute or concentrate spraying equipment. Apply the same amount of AC WHISTLER Miticide/Insecticide to the target crop whether by spraying by dilute or concentrate spraying methods.

Use the higher rate when pest pressure is high.

DO NOT apply more than once per season.

SITUATION	PEST	RATE/ha	WHP (days)
Cotton (Qld, NSW, WA ONLY) DO NOT apply to cotton under visible stress.	Carmine Spider Mite <i>(Tetranychus cinnabarinus)</i> Two Spotted Mite <i>(Tetranychus urticae)</i>	300 mL/ha	20
	Native budworm <i>(Helicoverpa punctigera)</i>	300 or 600 mL/ha	
Hops	Two Spotted Mite <i>(Tetranychus urticae)</i>	1L/ha	28
Strawberries	Two Spotted Mite <i>(Tetranychus urticae)</i>	100mL/100L If spray volume is less than 600L/ ha use a minimum of 600mL of this product /ha. DO NOT exceed 1200mL of this product /ha per application	3
Tomatoes -trellised or staked	Two Spotted Mite <i>(Tetranychus urticae)</i>	High volume spraying: 300mL/ha or 450mL/ha using concentrations of 60 to 90mL/ 100L	3
	Tomato Russet Mite <i>(Aculops lycopersici)</i>		
	Tobacco Leaf Miner (Potato moth) <i>(Phthorimaea operculella)</i>	High volume spraying: 600mL/ha using concentrations of 120mL/100L	

CRITICAL COMMENTS

DO NOT make more than two applications to cotton per season, regardless of pest being controlled.

To ensure adequate control:

1. Apply when the threshold number of mites has been reached for your area.
2. Apply when mite populations are low.

If the mite population is too high at time of application, satisfactory control may not be achieved.

Under high pest pressure a second application may be required 7 - 10 days after the initial application.

Thorough coverage is essential.

DO NOT make more than two applications to cotton per season, regardless of pest being controlled.

Use only when Lepton test kit results indicate that no greater than 10% *Helicoverpa armigera* are present. Use the higher rate alone or the lower rate with a suitable mixing partner.

Applications should target brown eggs and newly emerged larvae (neonates). Mixed size larval populations should be avoided.

Apply as a dilute spray in 1000L to 2000L water /ha, depending on crop size as pest pressure indicates.

DO NOT apply more than once per season.

Spray to wet all foliage to near the point of run-off. Ensure thorough coverage and penetration into plants. Best results are obtained if application is made at the first sign of mite appearance. When applied at this time one application may give good control. If mite numbers exceed 3 - 5 mites per leaflet, apply two applications 7 to 10 days apart. Re-apply if required but apply a maximum of 2 sprays of this product per season. If a further treatment is required, apply a product from a different chemical group.

Integrated Pest Control: see instructions in Apples and Pears section above. Refer to notes on resistance under General Instructions section of this label.

Spray to wet foliage to near the point of run-off. Ensure thorough coverage and penetration into plants. Best results are obtained if application is made before build up of mite numbers. Use the higher rate if mite numbers exceed 5 to 6 mites per compound leaf. Re-apply if required but apply a maximum of 2 sprays of this product per season. Allow at least 28 days between applications of this product. Refer to notes on resistance in the General Instructions section of this label.

Apply as per Two Spotted Mite. The lower rate will control Tomato Russet Mites not apparent at spraying. Use the higher rate if Tomato Russet Mite is present at spraying or is the main pest.

Wet foliage to the point of run-off. Thorough coverage and penetration into bushes is essential.

Apply at the first sign of pest activity. Re-apply as pest numbers indicate or every 7 to 10 days with a maximum of 5 applications to the crop. If mites are also a problem, DO NOT use more than 2 sprays per crop. Refer to notes on resistance in the General Instructions section of this label.

Note: Other IPM strategies to reduce leaf miner populations should be employed.

SITUATION	PEST	RATE/ha	WHP (days)
Tomatoes – field grown	Two Spotted Mite (<i>Tetranychus urticae</i>)	300 or 450mL/ha	3
	Tobacco leaf Miner (Potato moth) (<i>Phthorimaea operculella</i>)	600mL/ha	
	Tomato Russet Mite (<i>Aculops lycopersici</i>)	300 or 450mL/ha	
Ornamentals including Roses, Chrysanthemums, Carnations and indoor foliage plants	Two Spotted Mite (<i>Tetranychus urticae</i>)	50mL/100L to a maximum of 1.5L	Nil

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS:

COTTON:

DO NOT HARVEST FOR 20 DAYS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 20 DAYS AFTER APPLICATION.

CITRUS:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

HOPS:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

APPLES AND PEARS:

DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

DO NOT FEED TREATED PRODUCE TO LIVESTOCK FOR 14 DAYS AFTER APPLICATION.

CAPSICUMS, STRAWBERRIES AND TOMATOES:

DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

DO NOT FEED TREATED PRODUCE TO LIVESTOCK FOR 3 DAYS AFTER APPLICATION

ORNAMENTALS:

NOT REQUIRED WHEN USED AS DIRECTED

DO NOT GRAZE OR CUT FOR STOCK FOOD

CRITICAL COMMENTS

Apply in sufficient volume to obtain even coverage and penetration of plants. Use droppers to direct spray onto plants and away from the inter-row. Best results are obtained if application is made before build up of mite numbers. Use the higher rate if mite numbers exceed 5 mites per compound leaf. Re-apply if required but apply a maximum of 2 spray of this product per season. Allow at least 28 days between applications of this product. Refer to notes on resistance in the General Instructions section of this label.

Apply in sufficient volume to obtain even coverage and penetration of plants. Use droppers to direct spray onto plants away from the inter-row. Apply at the first sign of pest activity. Re-apply as pest numbers indicate or every 7 to 10 days with a maximum of 5 applications to the crop. If mites are also a problem, DO NOT use more than 2 AC WHISTLER Miticide/Insecticide sprays per crop. Refer to notes on resistance in the General Instructions section of this label.

Note: Other IPM strategies to reduce leafminer populations should be employed.

Apply as per Two Spotted Mite. The lower rate will control Tomato Russet Mites not apparent at spraying. Use the higher rate if Tomato Russet Mite is present at spraying or is the main pest.

Spray to wet foliage to near the point of run-off using at least 2000L water/ha (100L/500 m²). Thorough coverage and penetration into plants is essential. Preferably apply on first appearance of mites. When applied when pest numbers are low to moderate, one application will be sufficient to give effective control, however if mites are numerous, apply a second application 7 to 10 days later. DO NOT use overhead irrigation within 24 hours after application.

DO NOT use on Ferns or Shasta.

For ornamentals not listed on this label, small test applications to assess unexpected phytotoxicity should be made before spraying the whole crop.

DO NOT use more than 2 times per season.

Refer to notes on resistance in the General Instructions section of this label.

GENERAL INSTRUCTIONS:

This product is not systemic, but quickly moves into leaves following application. It is important that thorough coverage is achieved so that the maximum amount of product can be deposited on the leaf surface for uptake and subsequent ingestion by sucking mites. Product that is not absorbed by leaves is quickly degraded.

RESISTANCE

It is known that early development of mite resistance may occur if one particular miticide from one particular chemical group are continually used. For this reason it is recommended that this product be applied no more than twice in one season preferably only once per season. A second application is only recommended where mite pressure is very high. It is also recommended to alternate this product with miticides from a different chemical group. Consult your local supplier or technical adviser for further information on chemical groups.

INSECTICIDE RESISTANCE WARNING

GROUP 6 INSECTICIDE

AC WHISTLER MITICIDE/INSECTICIDE is a Group 6 miticide and insecticide. Some naturally-occurring insect biotypes resistant to this product and other Group 6 insecticides may exist through normal genetic variability in

any insect population. The resistant individuals can eventually dominate the insect population if the product or other Group 6 insecticides are used repeatedly. The effectiveness of the product on resistant individuals could be significantly reduced.

Since the occurrence of resistant individuals is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects. AC WHISTLER MITICIDE/INSECTICIDE may be subject to specific resistance management strategies. For further information contact your supplier, Axichem Pty Ltd representative or local agricultural department agronomist.

It is recommended that this product be applied no more than the following amount of times per season/crop and that it is not used consecutively except where a two-spray schedule is specifically recommended:

Crop	No of Sprays per Season
Cotton	No more than twice in one season and preferably only once per season. A second application is only recommended where mite pressure is very high.
Apples, Capsicums, Citrus, Hops, Pears	One spray per season.
Tomatoes	Two sprays per season if mites are present. Five sprays per season if mites are NOT present.
Strawberries	Two sprays per season.

AC WHISTLER MITICIDE/INSECTICIDE should not be applied in two consecutive seasons or crops without an unrelated chemical being used in between. Alternate this product with approved miticides from other chemical groups. For further information contact your local supplier, Axichem Pty Ltd representative or local department of agriculture agronomist.

CROP MONITORING

It is essential to carry out regular crop monitoring of mite levels (every 2 - 3 days for cotton and every 3 - 5 days for other crops) to ensure the product is applied at the correct timing.

MIXING INSTRUCTIONS FOR APPLES AND PEARS

To achieve 750mL this product + 5L Summer oil/ha apply spray at the following mixing rates:

	AC WHISTLER MITICIDE/INSECTICIDE	Summer Oil
(Quantity per 100L required)		
1000L/ha (minimum)	75mL	500mL
1500L/ha	50mL	335mL
2000L/ha	37.5mL	250mL
2500L/ha	30mL	200mL

MIXING

Partially fill the spray tank with water, add the required amount of AC WHISTLER MITICIDE/INSECTICIDE, and then add the remainder of the water. If oil is recommended then add this after the AC WHISTLER MITICIDE/INSECTICIDE is well mixed. A wetting agent is NOT required.

CROP SAFETY

A mixture of this product with summer oil has very occasionally caused slight fruit russetting on some pear varieties particularly Anjou and other sensitive varieties when used alone or when other products are applied sequentially. A very small amount of temporary apple fruit blemishing has been associated with low water volume applications. DO NOT apply AC WHISTLER MITICIDE/INSECTICIDE to apples or pears before or after applications of Delan* or Captan*. The label instructions on the summer oil label must be strictly followed. Conditions which may contribute to crop damage are:

- 1) unusually hot conditions present or expected within 24 hours of application
- 2) poor or slow drying conditions
- 3) application with equipment that may leave large droplets on fruit after application.

COMPATIBILITY

This product is compatible with most commonly used insecticides and fungicides. However all mixtures should be tested prior to mixing commercial quantities.

APPLICATION INSTRUCTIONS FOR COTTON

AC WHISTLER MITICIDE/INSECTICIDE may be applied by ground spraying equipment or by fixed wing aircraft. Apply in a minimum of 20L water per hectare. Ensure good coverage.

Ground application: Apply with inter-row droppers fitted with nozzles spraying towards the cottons rows. The inter-row nozzles should be level with or just below the canopy and spraying at right angles to the ground.

Aerial application (Fixed Wing): Apply in the cooler parts of the day or night when there is a reliable cross wind to assist with good penetration into the crop canopy. Preferably use aircraft fitted with micronair atomisers.

APPLICATION INSTRUCTIONS FOR CITRUS

To be effective this product requires thorough spray coverage. Ensure that equipment is properly calibrated to give an even distribution at the correct volume. The same quantity of product per hectare should be used when spraying either with dilute or concentrate spraying.

APPLICATION BY DILUTE SPRAYING

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set-up and operation may also need to be changed, as the crop grows.

APPLICATION BY CONCENTRATE SPRAYING

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only

- (i) Dilute spray volume as determined above: For example 1500L/ha
- (ii) Your chosen concentrate spray volume: For example 500L/ha
- (iii) The concentration factor in this example is: $3 \times$ (i.e. $1500\text{L} / 500\text{L} = 3$)
- (iv) If the dilute label rate is 10mL/100L, then the concentrate rate becomes 3×10 that is 30mL/100L of concentrate spray.
 - The chosen spray volume, amount of product per 100L of water and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTIONS

Re-entry period

Allow the spray to dry on the foliage before re-entry to treated areas.

DO NOT allow entry into treated areas in glasshouses for 24 hours after treatment. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and elbow-length PVC gloves.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

Drift Warning: Do NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK:

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Dangerous to fish and other aquatic organisms.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Studies indicate that when Abamectin comes into contact with soil it readily and tightly binds to the soil and becomes inactive over time.

STORAGE AND DISPOSAL:

Store below 30°C (Room temperature). Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture empty containers and deliver for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Poisonous if absorbed by skin contact or swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. For space applications, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles and a half face respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and

water. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26.

If swallowed, do NOT induce vomiting. Give a glass of water.

If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

If in eyes wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from Axichem Pty Ltd.

LABEL ELEMENTS

GHS Label Elements		
--------------------	---	---

SIGNAL WORD	DANGER
-------------	--------

Hazard Statement(s)

H301	Toxic if swallowed.
H315	Causes skin irritation.
H319	Causes serious eye irritation.
H331	Toxic if inhaled.
H335	May cause respiratory irritation.
H360Df	May damage the unborn child. Suspected of damaging fertility.
H373	May cause damage to organs through prolonged or repeated exposure.
H412	Harmful to aquatic life with long lasting effects.

Precautionary statement(s) Prevention

P201	Obtain special instructions before use.
P260	Do not breathe mist/vapours/spray.
P264	Wash all exposed external body areas thoroughly after handling.
P270	Do not eat, drink or smoke when using this product.
P271	Use only outdoors or in a well-ventilated area.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P273	Avoid release to the environment.

Precautionary statement(s) Response

P301+P310	IF SWALLOWED: Immediately call a POISON CENTRE/doctor/physician/first aider.
P308+P313	If exposed or concerned: Get medical advice/attention.
P330	Rinse mouth.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P311	Call a POISON CENTRE/doctor/physician/first aider.
P337+P313	If eye irritation persists: Get medical advice/attention.
P302+P352	IF ON SKIN: Wash with plenty of water.
P332+P313	If skin irritation occurs: Get medical advice/attention.
P362+P364	Take off contaminated clothing and wash it before reuse.

Precautionary statement(s) Storage

P403+P233	Store in a well-ventilated place. Keep container tightly closed.
P405	Store locked up.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

* Not a trademark of Axichem Pty Ltd