


DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as the n-octanoyl ester
SOLVENT: 616 g/L LIQUID HYDROCARBON

GROUP **6** HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and Lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions For Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions. Use only at recommended growth stages of crops and weeds. Do not use on crops other than recommended.

STORAGE AND DISPOSAL

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. Do NOT burn empty containers or product.

Refillable containers (1000 L only):

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia

13 11 26). If swallowed do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

LIMIT OF WARRANTY AND LIABILITY

AxiChem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of AxiChem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repackaging or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 81113/RV2024

Batch No:

Date of Manufacture:

IN A TRANSPORT EMERGENCY
 DIAL 000
 POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY
 CALL 1800 039 008
 ALL HOURS AUSTRALIA WIDE

AxiChem
 Pty Ltd

AXICHEM PTY LTD

A.B.N 35 131 628 594

9 PALINGS COURT, NERANG QLD 4211

PHONE: 07 5596 1736 www.axichem.com.au

PEE
 HERE

AxiChem

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply by mister machines.

DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.

DO NOT apply to diseased or frost affected crops or turf or if frosts are imminent.

DO NOT apply when rain is expected within 3 hours.

DO NOT apply to Lucerne crops or clover based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale: Including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only)	Amsinckia (<i>Amsinckia</i> spp.) Black Bindweed (Climbing buckwheat) (<i>Fallopia convolvulus</i>), Capeweed (<i>Arctotheca calendula</i>) Chamomile (<i>Matricaria matricarioides</i>), Common Peppergrass (<i>Lepidium africanum</i>), Corn Gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain sorrel (<i>Oxalis acetosella</i>), Purpose Calandrinia (<i>Calandrinia menziesii</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Three-horned bedstraw (<i>Gallium tricornutum</i>), Tree Hogweed (<i>Polygonum patulum</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 35mm in diameter	All States	1.4L/ha	CROP STAGE: Wheat, Barley, Cereal, Rye, Oats and Triticale: 3 leaf to full tillered (Zadoks Scale Z13-Z30). Linseed: 50 to 150mm high. Lucerne and Clover: After seedlings have one trifoliate leaf or before there is 150mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a minimum of 2-6 true leaves). APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian clover or Berseem clover.
	Mexican Poppy (<i>Argemone ochroleuca</i>), Three-cornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>), Turnip Weed (Giant Mustard) (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Mustard (<i>Sisymbrium</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed ^{##} (<i>Polygonum aviculare</i>), Paterson's Curse [*] (<i>Echium plantagineum</i>) (Salvation Jane)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter		2.0L/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only) (cont'd)	Fumitories* (<i>Fumaria</i> spp.)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Vic, SA, WA, ACT only	2.0L/ha	CAUTION: (cont'd) Linseed: DO NOT spray linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of Medics may result in unacceptable crop damage in some situations. #Effective control of Paterson's Curse and Fumitories will only be obtained under good growing conditions. ##Where Wireweed is growing under stress, it has been found to be less susceptible.
	Pheasant's Eye (<i>Adonis microcarpa</i>), Birds Eye (Carrot Weed) (<i>Cotula australis</i>)	Up to the 4 true leaf stage	SA only	700mL/ha	DO NOT use where undersown medics occur.
		Greater than the 4 true leaf stage		1.4L/ha	
	Fireweed (<i>Senecio</i> spp.)	Seedlings	Qld, NSW, ACT only	1.4L/ha	Apply during the Autumn-Winter period when weeds are young and actively growing. Not effective on mature plants.
		Early flowering stage		2.8L/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medics; Clover Seed Crops	Amsinckia (<i>Amsinckia</i> spp.), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvenses</i>), Corn Gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Hexham Scent (King Island Melilot) (<i>Melilotus indicus</i>), Horned Poppy (<i>Glaucium flavum</i>), Indian Hedge Mustard, (<i>Sisymbrium orientale</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mexican Poppy (<i>Argemone ochroleuca</i>), Mintweed (<i>Salvia reflexa</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>), Paterson's Curse (Salvation Jane) (<i>Echium plantagineum</i>), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Slender Thistle (<i>Carduus tenuiflorus</i>), Turnip Weed (Giant Mustard) (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>)	Up to the 5 leaf stage or when plants are no more than 50mm in diameter	WA only	750mL/ha PLUS 440mL Amine 625/ha OR PLUS 700mL of MCPA 500/ha	CROP STAGE: Cereals: 3 leaf to fully tillered. (Zadoks Scale Z13-30). Clover: Apply after clover seedlings have at least three trifoliate leaves APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover of Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with Lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Diclofop methyl.
	Common Peppergrass (<i>Lepidium africanum</i>). Tree Hogweed (<i>Polygonum patulum</i>)		Vic, SA only	1.4/ha PLUS 440mL Amine 625/ha OR PLUS 700mL of MCPA 500/ha	
	Black Bindweed (Climbing Buckwheat) (<i>Fallopia convolvulus</i>)	Up to the 8 leaf stage.	Vic, SA only		

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Oats, Barley, Triticale alone or with undersown Lucerne and clover		APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian clover or Berseem clover.			
	Common Sowthistle (<i>Sonchus oleraceus</i>)	Up to 4 leaves	Qld, Northern NSW	1.4 L/ha	Strictly for use on small weeds early in crop before shading out. CROP STAGE: Wheat, Barley, Oats and Triticale: 3 leaf to mid tillering (Z13-Z23). Use a minimum spray volume of 70 L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1 L/ha
		4 to 8 leaves		2.1 L/ha	CROP STAGE: Wheat, Barley, Oats and Triticale: 3 leaf to full tillering (Z13-Z30). This product will cause slight leaf burning of undersown Lucerne and clover.
Wheat, Barley, Oats and Triticale	Threecornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>), Wireweed (<i>Polygonum aviculare</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Tas, Sa, ACT only	1.4/ha PLUS 440mL 2,4-D Amine	CROP STAGE: 5 leaf to fully tillered (Zadoks scale Z15030): Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 200L spray per hectare on Black Bindweed when crop is greater than 300mm high. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with Lucerne, clover or medics unless some damage is acceptable.
	Fumitories (<i>Fumaria</i> spp.)		Qld, NSW, ACT, Vic only	625 / ha OR PLUS 700mL of MCPA	
	Wild Mustard (<i>Sisymbrium</i> spp.) Wild Turnip (<i>Brassica tournefortii</i>)		Qld, Vic only	500/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Oats and Triticale (cont'd)	Cleavers	2-4 stems + 1-3 whorls of leaves/ stem	Qld, Vic, NSW, ACT, Tas only	1.4/ha PLUS 440mL 2,4-D Amine 625 / ha OR PLUS 700mL of MCPA 500/ ha	CAUTION: (cont'd) DO NOT apply to medics (WA only). This product may be tank mixed with Diclofop methyl for control of Annual Rye Grass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using tank mix with Diclofop methyl. APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.
	Threecornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>) Wireweed (<i>Polygonum aviculare</i>)	4-5 leaf stage	WA only		
			Qld, Vic, NSW, ACT, SA, Tas only	2.1L/ha PLUS 400mL 2,4-D Amine 625 / ha OR PLUS 700mL of MCPA 500/ ha	
	Fumitories (<i>Fumaria</i> spp.)		Qld, NSW, Vic, SA, ACT only		
	Black Bindweed (Climbing Buckwheat) (<i>Fallopia convolvulus</i>), Chamomile (<i>Matricaria matricarioides</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain sorrel (<i>Oxalis acetosella</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>), Paterson's Curse (Salvation Jane) (<i>Echium plantagineum</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Three-horned bedstraw (<i>Gallium tricornutum</i>), Variegated Thistle (<i>Silybum marianum</i>)	5 to 8 leaf stage	All States		

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Fallow & Non Crop	Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>)	Up to 8 leaves	Qld, NSW, ACT only	1.4 L to 2.1 L/ha	Use the low rate for small weeds up to 6 leaves. Complete coverage is essential. Plants which are shaded by stubble will not be controlled. Application volumes above 50 L/ha are recommended. The addition of glyphosate will provide broader spectrum weed control where needed.
Pastures	Adonis (Pheasants Eye) Birds Eye (Carrot Weed)	Up to 4 leaves	SA only	700 mL	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. Do not apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application; as seedling mortality may occur and established plants may be damaged.
		More than 4 leaves		1.4 L	
	Double Gee, Wild Radish	Up to 4 leaves	WA only	1.5 L	
	Matricaria	Up to 3 leaves		750 mL	
	Fireweed	Seedlings	Qld, NSW, ACT only	1.4 L	
		Early flowering		2.8 L	
Pastures, Roadsides and Rights-of-Way	African Daisy (<i>Senecio pterophorus</i>)	Up to and including the flowering stage	SA only	300mL per 100L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. Spraymate Activator should be added at 125mL per 100 litres of spray. A follow-up treatment may be necessary for large, well-established plants. Ensure an overall spray coverage on weed seedlings.
	Boneseed/Bitou-bush (<i>Chrysanthemoides monilifera</i>)	Seedlings only	Vic, Tas only	160mL per 100L water	

Situation & Crop	Weeds Controlled	State	Rate	Critical Comments
Turf: Established Couch (including Queensland Blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue and Rye lawns	Bindy-eye (<i>Calotis hispidula</i>) Jo-jo (Onehunga) (<i>Soliva pterosperma</i>) Cudweed (<i>Gnaphalium</i> spp.)	All States	6.0L/500L water per ha High Volume: 300mL per 100L 60mL/100m ² per 15L knapsack	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

DO NOT GRAZE TREATED CROPS OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid instructions. Use only at recommended growth stages of crops and weeds. Do not use on crops other than recommended.

MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of water. Agitate again before spraying commences.

BOOM SPRAYER

Apply 50-200 litres (WA: 50 to 100 litres) of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense.

DO NOT use rain-drop or air-induction nozzles as coverage of weeds may not be adequate.

AERIAL APPLICATION

Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

EQUIPMENT MAINTENANCE

After use, wash equipment thoroughly and rise several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixes with Amine 625 or MCPA 500 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides or insecticides in wheat and barley only.

Diclofop methyl, Omethoate, Chlorpyrifos, dimethoate, 2,4-D Amine 625, MCPA 500 Herbicide, Chlorsulfuron,

Metsulfuron, Diflufenican, Glyphosate, Dicamba 500.

As formulations of other manufacturers products are beyond the control of Axichem Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANT WEEDS WARNING

GROUP	6	HERBICIDE
-------	---	-----------

AC Brag 200 Herbicide is a member of the nitriles group of herbicides. AC Brag 200 Herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management AC Brag 200 Herbicide is a Group 6 herbicide. Some naturally-occurring weed biotypes resistant to AC Brag 200 Herbicide and other Group 6 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 6 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Ltd. Accepts no liability for any losses that may result from the failure of AC Brag 200 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Note: Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

Refillable containers (1000 L only):

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before

eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.




FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

LABEL ELEMENTS

GHS Label Elements			
SIGNAL WORD	Danger		
Hazard Statement(s)			
H227	Combustible liquid		
H304	May be fatal if swallowed and enters airways.		
H312	Harmful in contact with skin.		
H317	May cause an allergic skin reaction.		
H319	Causes serious eye irritation.		
H332	Harmful if inhaled.		
H335	May cause respiratory irritation.		
H336	May cause drowsiness or dizziness.		
H361d	Suspected of damaging the unborn child.		
H373	May cause damage to organs through prolonged or repeated exposure.		
H410	Very toxic to aquatic life with long lasting effects.		
AUH066	Repeated exposure may cause skin dryness and cracking.		
Precautionary statement(s) Prevention			
P201	Obtain special instructions before use.		
P210	Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.		
P260	Do not breathe mist/vapours/spray.		
P271	Use only outdoor or in a well-ventilated area.		
P280	Wear protective gloves and protective clothing, eye protection and face protection.		

P273	Avoid release to the environment.
P264	Wash all exposed external body areas thoroughly after handling.
P272	Contaminated work clothing should not be allowed out of the workplace.

Precautionary statement(s) Response

P301+P310	IF SWALLOWED: Immediate call a POISON CENTRE/doctor/physician/first aider.
P331	Do NOT induce vomiting.
P308+P313	If exposed or concerned: Get medical advice/attention.
P370+P378	In case of fire: Use alcohol resistant foam or normal protein foam to extinguish.
P302+P352	IF ON SKIN: Wash with plenty of water and soap.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P312	Call a POISON CENTER/doctor/physician/first aider/if you feel unwell.
P333+P313	If skin irritation or rash occurs: Get medical advice/attention.
P337+P313	If eye irritation persists: Get medical advice/attention.
P362+P364	Take off contaminated clothing and wash it before reuse.
P391	Collect spillage.
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.

Precautionary statement(s) Storage

P405	Store locked up.
P403+P233	Store in a well-ventilated place. Keep container tightly closed.

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

AxiChem